

Need more money for college expenses?

The CLC Private LoanSM can get you up to
\$40,000 a year for college-related expenses.

Here's why the CLC Private LoanSM is a smart choice:

- ✓ Approved borrowers are sent a check within four business days
- ✓ Get \$1000 - \$40,000 each year
- ✓ Interest rates as low as prime + 0% (8.66% APR)
- ✓ Quick and easy approval process
- ✓ No payments until after graduating or leaving school

CLICK HERE

or call **800.311.9615.**

*We are available 24 hours
a day, 7 days a week.*

Roots and Radicals

Roger has a ladder that will extend to a length of 21 feet. For the ladder to be safe to climb on, it must be placed 7 feet away from the house. The roof is 20 feet above the ground. Will the ladder be able to reach the roof safely? If not, how far up will the ladder reach? (Leave your answer rounded to one decimal place.)



9-1 ■ Principal roots

Square root

In chapter 4, quadratic equations were solved by factoring, but many quadratic equations such as

$$x^2 - 7 = 0 \quad \text{or} \quad x^2 + 3x + 1 = 0$$

will not factor over the set of rational numbers. We need to be able to solve equations that involve a squared variable. Therefore, we want a process that is the inverse of squaring a number.

In chapter 1, we discussed how to square a number. Consider the following examples:

$$\text{If } x = 3, \text{ then } x^2 = (3)^2 = (3)(3) = 9$$

$$\text{If } x = -3, \text{ then } x^2 = (-3)^2 = (-3)(-3) = 9$$

Reversing the process, we ask:

$$\text{If } x^2 = 9, \text{ then what number is } x \text{ equal to?}$$

This inverse operation is called *finding the square root of a number*.

Definition

For every pair of real numbers a and b , if $a^2 = b$, then a is called a square root of b .

Concept

A square root of a number is one of two equal factors of the number.

From this discussion and the definition of square root, we can see that the answer to the question we asked,

If $x^2 = 9$, then what number is x equal to?

is 3 or -3 since $(3)^2 = 9$ and $(-3)^2 = 9$. To distinguish between the two square roots, we define the *principal square root* of a positive number to be positive. The $\sqrt{}$ symbol denotes the principal square root.

Thus, if $x = \sqrt{9}$, then $x = 3$ and we say 3 is the principal square root of 9.

$$\sqrt{9} = 3 \text{ (principal square root)}$$

The parts of the principal square root are



The entire expression is called a *radical* and is read “the principal square root of a .”

■ Example 9-1 A

Find the principal square root.

1. $\sqrt{16} = 4$, since $4 \cdot 4 = 4^2 = 16$.
2. $\sqrt{49} = 7$, since $7 \cdot 7 = 7^2 = 49$.
3. $\sqrt{25} = 5$, since $5 \cdot 5 = 5^2 = 25$.
4. $\sqrt{36} = 6$, since $6 \cdot 6 = 6^2 = 36$.
5. $\sqrt{0} = 0$, since $0 \cdot 0 = 0^2 = 0$.

Note In the examples, 0, 16, 49, 25, and 36 are called **perfect-square integers** because their square roots are integers.

Whenever we wish to express the negative value of the square root of a number, we use the symbol $-\sqrt{}$. For example, $-\sqrt{9}$ would indicate that we want the negative square root value. That is, $-\sqrt{9} = -3$.

■ Example 9-1 B

Find the indicated root.

- | | |
|----------------------|----------------------|
| 1. $-\sqrt{4} = -2$ | 3. $-\sqrt{25} = -5$ |
| 2. $-\sqrt{49} = -7$ | 4. $-\sqrt{36} = -6$ |

► **Quick check** Find the square root. $-\sqrt{16}$

Our first examples of finding the square root of a number have dealt only with perfect-square integers. We shall now try to find the $\sqrt{2}$. We could use 1.414 but when we square 1.414, we do not get 2 as an answer. We can show that no matter how many decimal places the answer is carried to, when the result is squared, it will be close to, but not equal, 2. The $\sqrt{2}$ is called an **irrational number** because it has the property that it cannot be expressed as a terminating or a repeating decimal number. Another number that is irrational is π , which is used in geometric formulas involving circles.

Whenever you work with irrational numbers in a problem, you may have to approximate the number to as many decimal places as are needed in the problem by using a calculator or a table of values.

■ Example 9-1 C

For the following irrational numbers, find the decimal approximation to three decimal places by using a calculator.

1. $\sqrt{3} \approx 1.732$

Note “ \approx ” is read “is approximately equal to” and is used when our answer is not exact. Square roots of integers that are not perfect squares will be irrational.

2. $-\sqrt{41} \approx -6.403$

3. $\sqrt{50} \approx 7.071$

► **Quick check** Find the decimal approximation to three decimal places by using a calculator. $\sqrt{48}$ and $-\sqrt{56}$

Not all real numbers have a rational or an irrational square root. Consider the following:

$$\sqrt{-4} = \text{what?}$$

We know that all real numbers are either positive, negative, or zero. If we square a real number, the product is never negative. Hence, there is no real number that when squared produces a negative answer. *The square root of a negative number does not exist in the set of real numbers.*

■ Example 9-1 D

Suppose an automotive engineer wishes to determine the diameter of the cylinder bore (D) required to produce H horsepower from N cylinders of an engine that is turning 1,000 rpm. The engineer will use the formula

$$D = \sqrt{\frac{H}{(0.4)N}}$$

What would be the bore diameter (in inches) required to produce 40 horsepower at 1,000 rpm from a 4-cylinder engine?

We substitute 40 for H and 4 for N .


$$D = \sqrt{\frac{(40)}{(0.4)(4)}} = \sqrt{\frac{40}{1.6}} = \sqrt{25} = 5$$

Therefore, we need a bore diameter of 5 inches.

Pythagorean Theorem

The following is an important property of right triangles called the **Pythagorean Theorem**.

In a right triangle, the square of the length of the hypotenuse (the side opposite the right angle) is equal to the sum of the squares of the lengths of the two legs (the sides that form the right angle). If c is the length of the hypotenuse and a and b are the lengths of the legs, this property can be stated as:



$$\begin{aligned} c^2 &= a^2 + b^2 \text{ or } c = \sqrt{a^2 + b^2} \\ \text{also as } a^2 &= c^2 - b^2 \text{ or } a = \sqrt{c^2 - b^2} \\ \text{and as } b^2 &= c^2 - a^2 \text{ or } b = \sqrt{c^2 - a^2} \end{aligned}$$

Example 9-1 E


Find the length of the hypotenuse of a right triangle whose legs are 6 centimeters and 8 centimeters.

We want to find c when $a = 6$ cm and $b = 8$ cm.

By the Pythagorean Theorem,

$$\begin{aligned} c &= \sqrt{a^2 + b^2} \\ &= \sqrt{6^2 + 8^2} && \text{Substitute} \\ &= \sqrt{36 + 64} && \text{Square the values} \\ &= \sqrt{100} = 10 \end{aligned}$$

Hence, $c = 10$ cm.



► **Quick check** Find the second leg of a right triangle whose hypotenuse has length 13 inches and whose first leg is 5 inches long.

n th roots

The concept of square root can be extended to find cube roots (third root of a number), fourth roots, fifth roots, and so on. A cube root is one of three equal factors of a number. The symbol that is used to express the principal cube root is $\sqrt[3]{}$. The 3 is called the **index** of the radical expression. The index denotes what root we are looking for. The principal fourth root would be indicated by $\sqrt[4]{}$. General notation for the principal n th root would be $\sqrt[n]{}$, where n is a natural number greater than 1.

The principal n th root

The principal n th root of number a , denoted by


$$\sqrt[n]{a}$$

is one of n equal factors such that $\sqrt[n]{a} = b$ and

$$\underbrace{b \cdot b \cdot b \cdots b}_{n \text{ factors}} = b^n = a$$

where n is a natural number greater than 1.

The parts of the principal n th root are



Note If there is no index associated with a radical symbol, it is understood to be 2.

If we exclude even roots of negative numbers, which do not exist in the set of real numbers, we can extend our idea of principal square root to all other roots by saying: **the principal n th root of a number, denoted by $\sqrt[n]{}$, has the same sign as the number itself.**

Example 9-1 F

Find the indicated root.

1. $\sqrt[4]{16} = 2$, since $2 \cdot 2 \cdot 2 \cdot 2 = 2^4 = 16$.
2. $\sqrt[3]{-27} = -3$, since $(-3)(-3)(-3) = (-3)^3 = -27$.
3. $\sqrt[3]{-125} = -5$, since $(-5)(-5)(-5) = (-5)^3 = -125$.
4. $\sqrt{-16}$ Does not exist in the set of real numbers.

► **Quick check** Find the cube root. $\sqrt[3]{-64}$

Mastery points**Can you**

- Find the principal square root of a perfect-square integer?
- Find the principal root of a number?
- Find the decimal approximation with a calculator for a root that is an irrational number?

Exercise 9-1

Find the indicated square root. See examples 9-1 A and B.

Example $-\sqrt{16}$ **Solution** $= -4$ Since $4 \cdot 4 = 16$, and we want the negative value of the square root

- | | | | | | |
|------------------|----------------|-----------------|-----------------|------------------|------------------|
| 1. $\sqrt{100}$ | 2. $\sqrt{36}$ | 3. $\sqrt{4}$ | 4. $\sqrt{64}$ | 5. $-\sqrt{144}$ | 6. $-\sqrt{81}$ |
| 7. $-\sqrt{121}$ | 8. $-\sqrt{4}$ | 9. $\sqrt{121}$ | 10. $\sqrt{81}$ | 11. $-\sqrt{16}$ | 12. $-\sqrt{64}$ |

Find the decimal approximation to three decimal places of the indicated square root by using a calculator. See example 9-1 C.


Examples $\sqrt{48}$ $-\sqrt{56}$ **Solutions** ≈ 6.928 ≈ -7.483

- | | | | | | |
|-----------------|-----------------|-----------------|-----------------|------------------|------------------|
| 13. $\sqrt{18}$ | 14. $\sqrt{24}$ | 15. $\sqrt{41}$ | 16. $\sqrt{47}$ | 17. $-\sqrt{52}$ | 18. $-\sqrt{10}$ |
|-----------------|-----------------|-----------------|-----------------|------------------|------------------|

Solve the following problems. See example 9-1 D.

- 19.** The current I (amperes) in a circuit is found by the formula $I = \sqrt{\frac{\text{watts}}{\text{ohms}}}$. What is the current of a circuit that has 3 ohms resistance and uses 1,728 watts?
- 20.** What is the current of a circuit that has 2 ohms resistance and uses 450 watts? (Refer to exercise 19.)

21. The slant height, S , of a right circular cone is found by the formula $S = \sqrt{r^2 + h^2}$, where r is the radius of the base and h is the height of the cone. What is the slant height of a right circular cone whose base radius is 5 units and whose height is 12 units?
22. What is the slant height of a right circular cone whose base radius is 3 units and whose height is 4 units? (Refer to exercise 21.)




In the following right triangles, find the length of the unknown side. See example 9–1 E.

Example Find the second leg of a right triangle whose hypotenuse has length 13 inches and whose first leg is 5 inches long.

Solution We want to find b given that $c = 13$ in. and $a = 5$ in. Using one of the forms of the theorem,

$$\begin{aligned} b &= \sqrt{c^2 - a^2} \\ &= \sqrt{(13)^2 - (5)^2} && \text{Substitute} \\ &= \sqrt{169 - 25} && \text{Square the values} \\ &= \sqrt{144} = 12 \end{aligned}$$

Hence $b = 12$ in.



- | | | |
|------------------------------|--------------------------------|-------------------------------|
| 23. $a = 3$ m, $b = 4$ m | 24. $a = 8$ ft, $c = 10$ ft | 25. $a = 12$ in., $b = 5$ in. |
| 26. $a = 15$ cm, $b = 8$ cm | 27. $a = 6$ yd, $c = 10$ yd | 28. $b = 16$ m, $c = 20$ m |
| 29. $a = 12$ mm, $b = 16$ mm | 30. $a = 10$ in., $b = 24$ in. | |

Find the indicated root. See example 9–1 F.

Example $\sqrt[3]{-64}$

Solution $= -4$ Since $(-4)^3 = -64$

- | | | | | | |
|----------------------|----------------------|---------------------|----------------------|--------------------|---------------------|
| 31. $\sqrt[3]{8}$ | 32. $\sqrt[3]{27}$ | 33. $\sqrt[3]{125}$ | 34. $\sqrt[3]{64}$ | 35. $\sqrt[3]{-8}$ | 36. $\sqrt[3]{-64}$ |
| 37. $\sqrt[4]{81}$ | 38. $\sqrt[4]{16}$ | 39. $-\sqrt[4]{81}$ | 40. $-\sqrt[4]{625}$ | 41. $\sqrt[5]{32}$ | 42. $\sqrt[6]{64}$ |
| 43. $-\sqrt[5]{243}$ | 44. $\sqrt[5]{-243}$ | 45. $\sqrt[10]{1}$ | 46. $\sqrt[14]{1}$ | 47. $\sqrt[9]{-1}$ | 48. $\sqrt[15]{-1}$ |

Solve the following problems. See example 9–1 D.

49. The formula for finding the length of an edge, e , of a cube when the volume, v , is known is $e = \sqrt[3]{v}$. What is the length of the edge of a cube whose volume is 729 cubic units?
50. What is the length of the edge of a cube whose volume is 216 cubic units? (Refer to exercise 49.)

Review exercises

Rewrite the following numbers in prime factor form. See section 1–1.

- | | | | | | | | |
|------|-------|------|-------|-------|-------|-------|-------|
| 1. 9 | 2. 12 | 3. 8 | 4. 40 | 5. 50 | 6. 81 | 7. 64 | 8. 16 |
|------|-------|------|-------|-------|-------|-------|-------|

9-2 ■ Product property for radicals

Multiplying square roots

In this section, we are going to develop properties for simplifying radicals. Consider the example

$$\sqrt{4} \cdot \sqrt{25} = 2 \cdot 5 = 10$$

We also observe that

$$\sqrt{4 \cdot 25} = \sqrt{100} = 10$$

From our example, we can conclude that

$$\sqrt{4} \cdot \sqrt{25} = \sqrt{4 \cdot 25}$$

We now can generalize the product property for square roots.

Product property for square roots

For all nonnegative real numbers a and b ,

$$\sqrt{a} \cdot \sqrt{b} = \sqrt{ab}$$

Concept

The product of two square roots is equal to the square root of their product.

■ Example 9-2 A

Perform the indicated operations. Assume that all variables represent nonnegative real numbers.

$$1. \sqrt{3}\sqrt{5} = \sqrt{3 \cdot 5} = \sqrt{15}$$

$$2. \sqrt{6}\sqrt{7} = \sqrt{6 \cdot 7} = \sqrt{42}$$

$$3. \sqrt{3}\sqrt{a} = \sqrt{3a}$$

$$4. \sqrt{x}\sqrt{y}\sqrt{z} = \sqrt{xyz}$$

Simplifying principal square roots

An important use of our product property is in simplifying radicals. Consider the following example:

Since 12 can be factored into $4 \cdot 3$, by our product property, we can write

$$\sqrt{12} = \sqrt{4 \cdot 3} = \sqrt{4}\sqrt{3} = 2\sqrt{3}$$

We are able to simplify the radical because the radicand contains a perfect-square integer factor, 4. In our example, $2\sqrt{3}$ is called the *simplified form* of $\sqrt{12}$.

Simplifying the principal square root

1. If the radicand is a perfect-square integer, write the corresponding square root.
2. If possible, factor the radicand so that at least one factor is a perfect-square integer. Write the corresponding square root as a coefficient of the radical.
3. The square root is in simplest form when the radicand has no perfect-square integer factors other than 1.

The following property is used when changing radicals involving variables in the radicand to simplest form.

$\sqrt{a^2}$ property

If a is any nonnegative real number, then

$$\sqrt{a^2} = a$$

■ Example 9-2 B

Simplify the following expressions. Assume that all variables represent nonnegative real numbers.

- $$\begin{aligned}\sqrt{50} &= \sqrt{25 \cdot 2} && \text{Factor having a perfect-square integer} \\ &= \sqrt{25} \sqrt{2} && \text{Product property} \\ &= 5\sqrt{2} && \sqrt{25} = 5\end{aligned}$$
- $$\sqrt{28} = \sqrt{4 \cdot 7} = \sqrt{4} \sqrt{7} = 2\sqrt{7}$$
- $$\sqrt{9a} = \sqrt{9 \cdot a} = \sqrt{9} \sqrt{a} = 3\sqrt{a}$$
- $$\sqrt{a^3} = \sqrt{a^2 \cdot a} = \sqrt{a^2} \sqrt{a} = a\sqrt{a}$$
- $$\sqrt{a^3 b^4} = \sqrt{a^2 \cdot a \cdot b^2 \cdot b^2} = \sqrt{a^2} \sqrt{a} \sqrt{b^2} \sqrt{b^2} = a\sqrt{abb} = ab^2\sqrt{a}$$
- $$\sqrt{a^2 + b^2}$$
 will not simplify because we are not able to factor the radicand.
The radicand must always be in a factored form before we can simplify.

Note $\sqrt{a^2 + b^2} \neq \sqrt{a^2} + \sqrt{b^2}$. For example, $\sqrt{3^2 + 4^2} = \sqrt{9 + 16} \neq \sqrt{9} + \sqrt{16}$. Since $\sqrt{9 + 16} = \sqrt{25} = 5$, whereas $\sqrt{9} + \sqrt{16} = 3 + 4 = 7$.

- $$\sqrt{2} \sqrt{6} = \sqrt{2 \cdot 6} = \sqrt{12} = \sqrt{4 \cdot 3} = \sqrt{4} \cdot \sqrt{3} = 2\sqrt{3}$$
- $$\sqrt{10} \sqrt{5} = \sqrt{10 \cdot 5} = \sqrt{50} = \sqrt{25 \cdot 2} = \sqrt{25} \sqrt{2} = 5\sqrt{2}$$
- $$\begin{aligned}\sqrt{14x} \sqrt{2x} &= \sqrt{14x \cdot 2x} = \sqrt{28x^2} = \sqrt{4 \cdot 7 \cdot x^2} = \sqrt{4} \sqrt{7} \sqrt{x^2} \\ &= 2\sqrt{7}x = 2x\sqrt{7}\end{aligned}$$

► **Quick check** Simplify. $\sqrt{8a^4}$ and $\sqrt{3a}\sqrt{6a}$

Multiplying n th roots

In section 9-1, we observed that even roots of negative numbers do not exist in the set of real numbers. Therefore, **we will consider all variables to be representing nonnegative real numbers whenever the index of the radical is even.**

Our product property for square roots can be extended to radicals with any index.

Product property for radicals

$$\sqrt[n]{a} \sqrt[n]{b} = \sqrt[n]{ab}$$

Concept

When we multiply two radicals *having the same index*, we multiply the radicands and put the product under a radical symbol with the common index.

■ Example 9-2 C

Perform the indicated operations.

- $\sqrt[3]{3}\sqrt[3]{2} = \sqrt[3]{3 \cdot 2} = \sqrt[3]{6}$
- $\sqrt[5]{7}\sqrt[5]{9} = \sqrt[5]{7 \cdot 9} = \sqrt[5]{63}$
- $\sqrt[3]{3}\sqrt[4]{5}$ These radicals cannot be multiplied together in this form since they do not have the same index. ■

Simplifying principal n th roots

We simplify n th roots, where n is greater than 2, as we did square roots. As long as the radicand can be factored so that one or more factors is a

- perfect cube when the index is 3,
- perfect fourth-power when the index is 4,
- perfect fifth-power when the index is 5, and so on,

the radical can be simplified. To do this, we use the property

$$\sqrt[n]{a^n} = a$$

where a is a nonnegative real number.

■ Example 9-2 D

Simplify the following radical expressions. Assume that all variables represent nonnegative real numbers.

- $$\begin{aligned}\sqrt[3]{81} &= \sqrt[3]{27 \cdot 3} && \text{Factor having a perfect-cube integer} \\ &= \sqrt[3]{27} \sqrt[3]{3} && \text{Product property} \\ &= 3\sqrt[3]{3} && \sqrt[3]{27} = 3\end{aligned}$$
- $\sqrt[4]{32} = \sqrt[4]{16 \cdot 2} = \sqrt[4]{16} \sqrt[4]{2} = 2\sqrt[4]{2}$
- $\sqrt[3]{x^5} = \sqrt[3]{x^3 \cdot x^2} = \sqrt[3]{x^3} \sqrt[3]{x^2} = x\sqrt[3]{x^2}$
- $\sqrt[5]{y^{10}} = \sqrt[5]{y^5 \cdot y^5} = \sqrt[5]{y^5} \sqrt[5]{y^5} = y \cdot y = y^2$

Note In example 4, the exponent 10 is evenly divisible by the index 5, and the radical is eliminated. When the exponent of a factor is evenly divisible by the index, that factor will no longer remain under the radical symbol.

- $\sqrt[4]{x^7 y^4} = \sqrt[4]{x^4 \cdot x^3 \cdot y^4} = \sqrt[4]{x^4} \sqrt[4]{x^3} \sqrt[4]{y^4} = x \sqrt[4]{x^3} y = xy \sqrt[4]{x^3}$
- $\sqrt[3]{a^7 b^2} = \sqrt[3]{a^3 a^3 a b^2} = \sqrt[3]{a^3} \sqrt[3]{a^3} \sqrt[3]{a b^2} = a \cdot a \sqrt[3]{a b^2} = a^2 \sqrt[3]{a b^2}$

Note No simplification relative to b is possible because the exponent of b is less than the value of the index.

$$7. \sqrt[3]{54x^3y^5} = \sqrt[3]{27 \cdot 2 \cdot x^3 \cdot y^3 \cdot y^2} = \sqrt[3]{27} \sqrt[3]{x^3} \sqrt[3]{y^3} \sqrt[3]{2y^2} = 3xy \sqrt[3]{2y^2}$$

► **Quick check** Simplify. $\sqrt[3]{8a^4}$ ■

Observe from the preceding examples that we can simplify a radical if the radicand has a factor(s) whose exponent is equal to or greater than the index.

Example 9-2 E

Perform the indicated multiplication and simplify. Assume that all variables represent nonnegative real numbers.

$$\begin{aligned} 1. \sqrt[3]{a^2} \sqrt[3]{a} &= \sqrt[3]{a^2 \cdot a} && \text{Product property} \\ &= \sqrt[3]{a^3} && \text{Multiply like bases} \\ &= a && \text{Perfect cube} \end{aligned}$$

$$\begin{aligned} 2. \sqrt[4]{8a^2} \sqrt[4]{4a^3} &= \sqrt[4]{8a^2 \cdot 4a^3} = \sqrt[4]{32a^5} = \sqrt[4]{16 \cdot 2 \cdot a^4 \cdot a} \\ &= \sqrt[4]{16} \sqrt[4]{a^4} \sqrt[4]{2a} = 2a\sqrt[4]{2a} \end{aligned}$$

► **Quick check** Simplify. $\sqrt[3]{9a^2} \sqrt[3]{3a^2}$

Note A very common error in problems involving radicals is to forget to carry along the correct index for the radical symbol.

Mastery points

Can you

- Multiply radicals having the same index?
- Simplify radicals?

Exercise 9-2

Perform any indicated operations and simplify. Assume that all variables represent nonnegative real numbers. See examples 9-2 A and B.

Examples $\sqrt{8a^4}$

$$\begin{aligned} \text{Solutions } \sqrt{8a^4} &= \sqrt{4 \cdot 2a^2a^2} && \text{Factor having perfect squares} \\ &= \sqrt{4} \sqrt{a^2} \sqrt{a^2} \sqrt{2} && \text{Product property} \\ &= 2a\sqrt{2} && \sqrt{4} = 2, \sqrt{a^2} = a \\ &= 2a^2\sqrt{2} && \text{Multiply} \end{aligned}$$

$\sqrt{3a} \sqrt{6a}$

$$\begin{aligned} &= \sqrt{3a \cdot 6a} && \text{Product property} \\ &= \sqrt{18a^2} && \text{Multiply} \\ &= \sqrt{9 \cdot 2 \cdot a^2} && \text{Factor having perfect squares} \\ &= \sqrt{9} \sqrt{2} \sqrt{a^2} && \text{Product property} \\ &= 3\sqrt{2} a && \sqrt{9} = 3, \sqrt{a^2} = a \\ &= 3a\sqrt{2} && \text{Multiply} \end{aligned}$$

- | | | | | | |
|--------------------------|---------------------------|----------------------------|-----------------------------|--------------------------|-----------------|
| 1. $\sqrt{16}$ | 2. $\sqrt{63}$ | 3. $\sqrt{20}$ | 4. $\sqrt{75}$ | 5. $\sqrt{45}$ | 6. $\sqrt{48}$ |
| 7. $\sqrt{32}$ | 8. $\sqrt{27}$ | 9. $\sqrt{80}$ | 10. $\sqrt{54}$ | 11. $\sqrt{98}$ | 12. $\sqrt{96}$ |
| 13. $\sqrt{a^7}$ | 14. $\sqrt{a^5}$ | 15. $\sqrt{4a^2b^3}$ | 16. $\sqrt{9ab^4c^3}$ | 17. $\sqrt{27a^3b^5}$ | |
| 18. $\sqrt{24x^5yz^3}$ | 19. $\sqrt{6}\sqrt{3}$ | 20. $\sqrt{27}\sqrt{6}$ | 21. $\sqrt{15}\sqrt{15}$ | 22. $\sqrt{11}\sqrt{11}$ | |
| 23. $\sqrt{6}\sqrt{10}$ | 24. $\sqrt{18}\sqrt{24}$ | 25. $\sqrt{25}\sqrt{15}$ | 26. $\sqrt{20}\sqrt{20}$ | 27. $\sqrt{5}\sqrt{15}$ | |
| 28. $\sqrt{2a}\sqrt{3a}$ | 29. $\sqrt{5x}\sqrt{15x}$ | 30. $\sqrt{6x}\sqrt{14xy}$ | 31. $\sqrt{2a}\sqrt{24b^2}$ | | |

32. A square-shaped television picture tube has a surface area of 121 square inches. What is the length of the side of the tube? (*Hint:* Area of a square is found by squaring the length of a side. $A = s^2$.)
33. A room in the shape of a square is 169 square feet. What is the length of a side? (See exercise 32.)
34. The formula for approximating the velocity V in miles per hour of a car based on the length of its skid marks S (in feet) on wet pavement is given by $V = 2\sqrt{3S}$. If the skid marks are 75 feet long, what was the velocity?
35. The formula for approximating the velocity V in miles per hour of a car based on the length of its skid marks S (in feet) on dry pavement is given by $V = 2\sqrt{6S}$. If the skid marks are 24 feet long, what was the velocity?

Perform any indicated operations and simplify. Assume that all variables represent positive real numbers. See examples 9-2 D and E.

Examples $\sqrt[3]{8a^4}$

Solutions $= \sqrt[3]{8 \cdot a^3 \cdot a}$ Factor having perfect cubes
 $= \sqrt[3]{8} \sqrt[3]{a^3} \sqrt[3]{a}$ Product property
 $= 2a \sqrt[3]{a}$ $\sqrt[3]{8} = 2, \sqrt[3]{a^3} = a$

$$\sqrt[3]{9a^2} \sqrt[3]{3a^2}$$

$= \sqrt[3]{9a^2 \cdot 3a^2}$ Product property
 $= \sqrt[3]{27a^4}$ Multiply radicands
 $= \sqrt[3]{27 \cdot a^3 \cdot a}$ Factor having perfect cubes
 $= \sqrt[3]{27} \sqrt[3]{a^3} \sqrt[3]{a}$ Product property
 $= 3a \sqrt[3]{a}$ $\sqrt[3]{27} = 3, \sqrt[3]{a^3} = a$

36. $\sqrt[3]{48}$ 37. $\sqrt[5]{64}$ 38. $\sqrt[4]{32}$ 39. $\sqrt[3]{24}$ 40. $\sqrt[5]{a^7}$
 41. $\sqrt[3]{b^8}$ 42. $\sqrt[3]{x^9}$ 43. $\sqrt[5]{y^{15}}$ 44. $\sqrt[3]{a^{12}}$ 45. $\sqrt[3]{4a^2b^3}$
 46. $\sqrt[3]{8r^2s^8}$ 47. $\sqrt[3]{16a^4b^5}$ 48. $\sqrt[5]{64x^{10}y^{14}}$ 49. $\sqrt[3]{81a^5b^{11}}$ 50. $\sqrt[3]{a^2} \sqrt[3]{a}$
 51. $\sqrt[3]{b^2} \sqrt[3]{b^2}$ 52. $\sqrt[5]{b^4} \sqrt[5]{b^3}$ 53. $\sqrt[5]{a} \sqrt[5]{a^4}$ 54. $\sqrt[3]{5a^2b} \sqrt[3]{75a^2b^2}$
 55. $\sqrt[3]{3ab^2} \sqrt[3]{18a^2b^2}$ 56. $\sqrt[4]{8a^3b} \sqrt[4]{4a^2b^2}$ 57. $\sqrt[4]{27a^2b^3} \sqrt[4]{9ab}$ 58. $\sqrt[3]{25x^5y^7} \sqrt[3]{15xy^3}$
 59. $\sqrt[3]{16a^{11}b^4} \sqrt[3]{12a^4b^6}$ 60. $\sqrt[4]{8xy} \sqrt[4]{4x^3y^3}$
61. The moment of inertia for a rectangle is given by the formula $I = \frac{bh^3}{12}$. If we know the values of I and b , we can solve for h as follows:
 $h = \sqrt[3]{\frac{12I}{b}}$. Find h if $I = 2 \text{ in.}^4$ and $b = 3 \text{ in.}$
62. Use exercise 61 to find h if $I = 27 \text{ in.}^4$ and $b = 4 \text{ in.}$
63. The moment of inertia for a circle is given by the formula $I = \frac{\pi r^4}{4}$. If we know the value of I , we can solve for r as follows: $r = \sqrt[4]{\frac{4I}{\pi}}$. Find r if $I = 12.56 \text{ in.}^4$ and we use 3.14 for π .
64. Use exercise 63 to find r if $I = 63.585 \text{ in.}^4$

Review exercises

Reduce the following fractions and rational expressions to lowest terms. Assume that no variable is equal to zero. See sections 1-1, 3-3, and 5-2.

1. $\frac{49}{56}$

2. $\frac{16x^3y^2}{-4xy}$

3. $\frac{2y^2 - 50}{y^2 - 4y - 5}$

Perform the indicated operations and simplify. Assume that all variables represent nonnegative real numbers. See sections 9-1 and 9-2.

4. $\sqrt{5}\sqrt{5}$

5. $\sqrt{3}\sqrt{3}$

6. $\sqrt{2}\sqrt{8}$

7. $\sqrt{12}\sqrt{3}$

8. $\sqrt{x}\sqrt{x}$

9-3 ■ Quotient property for radicals

The square root of a fraction

The following example will help us develop a property for division involving radicals.

$$\sqrt{\frac{4}{9}} = \sqrt{\left(\frac{2}{3}\right)^2} = \frac{2}{3}$$

We also observe that

$$\frac{\sqrt{4}}{\sqrt{9}} = \frac{2}{3}$$

From our example, we can conclude that

$$\sqrt{\frac{4}{9}} = \frac{\sqrt{4}}{\sqrt{9}}$$

We can now generalize this idea.

Quotient property for square roots

For any nonnegative real numbers a and b , where $b \neq 0$,

$$\sqrt{\frac{a}{b}} = \frac{\sqrt{a}}{\sqrt{b}}$$

Concept

The square root of a fraction can be written as the square root of the numerator divided by the square root of the denominator.

Example 9-3 A

Simplify the following expressions. Assume that all variables represent positive real numbers.

$$\begin{aligned} 1. \quad \sqrt{\frac{16}{25}} &= \frac{\sqrt{16}}{\sqrt{25}} \\ &= \frac{4}{5} \end{aligned}$$

Rewrite as the square root of the numerator over the square root of the denominator and simplify

$$2. \quad \sqrt{\frac{36}{49}} = \frac{\sqrt{36}}{\sqrt{49}} = \frac{6}{7}$$

$$3. \quad \sqrt{\frac{81}{100}} = \frac{\sqrt{81}}{\sqrt{100}} = \frac{9}{10}$$

$$4. \quad \sqrt{\frac{x^4}{64}} = \frac{\sqrt{x^4}}{\sqrt{64}} = \frac{x^2}{8}$$

$$5. \quad \sqrt{\frac{x^3}{y^4}} = \frac{\sqrt{x^3}}{\sqrt{y^4}} = \frac{x\sqrt{x}}{y^2}$$

► **Quick check** Simplify. $\sqrt{\frac{16}{49}}$ and $\sqrt{\frac{a^2}{b}}$

Rationalizing the denominator

When simplifying and evaluating radical expressions containing a radical in the denominator, it is easier if we can eliminate the radical in the denominator. For example,

$$\sqrt{\frac{4}{5}} = \frac{\sqrt{4}}{\sqrt{5}} = \frac{2}{\sqrt{5}}$$

Since $\sqrt{5} \cdot \sqrt{5} = 5$, we can eliminate the radical $\sqrt{5}$ in the denominator by multiplying the numerator and the denominator of the fraction by $\sqrt{5}$.

$$\frac{2}{\sqrt{5}} \cdot \frac{\sqrt{5}}{\sqrt{5}} = \frac{2\sqrt{5}}{\sqrt{25}} = \frac{2\sqrt{5}}{5}$$

The process of changing the denominator from a radical to a rational number is called **rationalizing the denominator**.

Rationalizing the denominator

1. Multiply the numerator and the denominator by the square root that is in the denominator. The radicand in the denominator will be a perfect-square integer.
2. Simplify the radical expressions in the numerator and the denominator.
3. Reduce the resulting fraction if possible.

■ Example 9-3 B

Simplify the following expressions. Leave no radicals in the denominator. Assume that all variables represent positive real numbers.

$$1. \frac{5}{\sqrt{7}} = \frac{5}{\sqrt{7}} \cdot \frac{\sqrt{7}}{\sqrt{7}} \quad \text{Multiply numerator and denominator by } \sqrt{7}$$

$$= \frac{5\sqrt{7}}{\sqrt{49}} \quad \text{Multiply in numerator and denominator}$$

$$= \frac{5\sqrt{7}}{7} \quad \sqrt{49} = 7$$

$$2. \frac{4}{\sqrt{6}} = \frac{4}{\sqrt{6}} \cdot \frac{\sqrt{6}}{\sqrt{6}} = \frac{4\sqrt{6}}{\sqrt{36}} = \frac{4\sqrt{6}}{6} = \frac{2\sqrt{6}}{3}$$

Note In example 2, we were able to reduce the fraction as a final step. Always check to see that the answer is in reduced form.

$$3. \frac{a}{\sqrt{a}} = \frac{a}{\sqrt{a}} \cdot \frac{\sqrt{a}}{\sqrt{a}} = \frac{a\sqrt{a}}{\sqrt{a^2}} = \frac{a\sqrt{a}}{a} = \sqrt{a}$$

$$4. \sqrt{\frac{a^3}{b}} = \frac{\sqrt{a^3}}{\sqrt{b}} = \frac{a\sqrt{a}}{\sqrt{b}} \cdot \frac{\sqrt{b}}{\sqrt{b}} = \frac{a\sqrt{ab}}{\sqrt{b^2}} = \frac{a\sqrt{ab}}{b}$$

► **Quick check** Simplify. $\frac{6}{\sqrt{6}}$

The following is a summary of the conditions necessary for a radical expression to be in **simplest form**, also called **standard form**.

1. The radicand contains no factors that can be written with an exponent greater than or equal to the index. ($\sqrt{a^3}$ violates this.)
2. The radicand contains no fractions. ($\sqrt{\frac{a}{b}}$ violates this.)
3. No radicals appear in the denominator. ($\frac{1}{\sqrt{a}}$ violates this.)

The n th root of a fraction

Our quotient property for square roots can be extended to radicals with any index.

Quotient property for radicals

$$\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}} \quad (b \neq 0)$$

Concept

The n th root of a fraction can be written as the n th root of the numerator divided by the n th root of the denominator.

Example 9-3 C

Simplify the following expressions. Assume that all variables represent positive real numbers.

$$\begin{aligned} 1. \quad \sqrt[3]{\frac{8}{27}} &= \frac{\sqrt[3]{8}}{\sqrt[3]{27}} \\ &= \frac{2}{3} \end{aligned}$$

Rewrite as the cube root of the numerator over the cube root of the denominator and simplify

$$2. \quad \sqrt[5]{\frac{32}{a^5}} = \frac{\sqrt[5]{32}}{\sqrt[5]{a^5}} = \frac{2}{a}$$

$$3. \quad \sqrt[3]{\frac{x^5}{y^6}} = \frac{\sqrt[3]{x^5}}{\sqrt[3]{y^6}} = \frac{x\sqrt[3]{x^2}}{y^2}$$

► **Quick check** Simplify. $\sqrt[3]{\frac{1}{27}}$ and $\sqrt[3]{\frac{8a^5}{b^3}}$

Rationalizing the denominator (n th root)

The following example will help us develop a general rule for rationalizing a denominator that has a single term.

$$\sqrt[3]{\frac{1}{a}} = \frac{\sqrt[3]{1}}{\sqrt[3]{a}} = \frac{1}{\sqrt[3]{a}}$$

At this point, a radical still remains in the denominator. We must now determine what we can do to the fraction to remove the radical from the denominator.

Observations:

1. We can multiply the numerator and the denominator by the same number and form equivalent fractions.
2. If we multiply by a radical, the indices must be the same to carry out the multiplication.
3. To bring a factor out from under the radical symbol and not leave any of the factor behind, the index must divide evenly into the exponent.

With these observations in mind, we rationalize the fraction as follows:

$$\begin{aligned} &= \frac{1}{\sqrt[3]{a}} \cdot \frac{\sqrt[3]{a^2}}{\sqrt[3]{a^2}} \\ &= \frac{1}{\sqrt[3]{a}} \cdot \frac{\sqrt[3]{a^2}}{\sqrt[3]{a^2}} \end{aligned}$$

Indices are the same

Multiply numerator and denominator by the same number

$$\begin{aligned}
 &= \frac{\sqrt[3]{a^2}}{\sqrt[3]{a^3}} \\
 &= \frac{\sqrt[3]{a^2}}{a}
 \end{aligned}$$

The sum of the exponents of a in the denominator is equal to the index

The index divides evenly into the exponent, the radical is eliminated

To rationalize an n th root denominator

1. Multiply the numerator and the denominator by a radical with the same index as the radical that we wish to eliminate from the denominator.
2. The exponent of each factor under the radical must be such that when we add it to the original exponent of the factor under the radical in the denominator, the sum will be equal to or divisible by the index of the radical.
3. Carry out the multiplication and reduce the fraction if possible.

■ Example 9-3 D

Simplify the following expressions. Leave no radicals in the denominator. Assume that all variables represent positive real numbers.

$$\begin{aligned}
 1. \quad \frac{1}{\sqrt[3]{7}} &= \frac{1}{\sqrt[3]{7}} \cdot \frac{\sqrt[3]{7^2}}{\sqrt[3]{7^2}} && \text{Multiply numerator and denominator by } \sqrt[3]{7^2} \\
 &= \frac{\sqrt[3]{7^2}}{\sqrt[3]{7^3}} && \text{Multiply in numerator and denominator } (\sqrt[3]{7} \sqrt[3]{7^2} = \sqrt[3]{7^3}) \\
 &= \frac{\sqrt[3]{7^2}}{7} && \sqrt[3]{7^3} = 7 \\
 &= \frac{\sqrt[3]{49}}{7} && 7^2 = 49
 \end{aligned}$$

$$2. \quad \frac{a}{\sqrt[5]{b^2}} = \frac{a}{\sqrt[5]{b^2}} \cdot \frac{\sqrt[5]{b^3}}{\sqrt[5]{b^3}} = \frac{a\sqrt[5]{b^3}}{\sqrt[5]{b^5}} = \frac{a\sqrt[5]{b^3}}{b}$$

$$3. \quad \frac{x}{\sqrt[4]{x}} = \frac{x}{\sqrt[4]{x}} \cdot \frac{\sqrt[4]{x^3}}{\sqrt[4]{x^3}} = \frac{x\sqrt[4]{x^3}}{\sqrt[4]{x^4}} = \frac{x\sqrt[4]{x^3}}{x} = \sqrt[4]{x^3}$$

$$4. \quad \frac{1}{\sqrt[5]{a^2b}} = \frac{1}{\sqrt[5]{a^2b}} \cdot \frac{\sqrt[5]{a^3b^4}}{\sqrt[5]{a^3b^4}} = \frac{\sqrt[5]{a^3b^4}}{\sqrt[5]{a^5b^5}} = \frac{\sqrt[5]{a^3b^4}}{ab}$$

► **Quick check** Simplify. $\frac{1}{\sqrt[5]{b^2}}$

Mastery points

Can you

- Simplify radicals containing fractions?
- Rationalize denominators?

Exercise 9-3

Simplify the following expressions. Leave no radicals in the denominator. Assume that all variables represent positive real numbers. See examples 9-3 A and B.

Examples $\sqrt{\frac{16}{49}}$

Solutions $= \frac{\sqrt{16}}{\sqrt{49}}$
 $= \frac{4}{7}$

Rewrite as the square root of the numerator over the square root of the denominator and simplify

$$\sqrt{\frac{a^2}{b}}$$

$$= \frac{\sqrt{a^2}}{\sqrt{b}}$$

$$= \frac{a}{\sqrt{b} \sqrt{b}}$$

$$= \frac{a\sqrt{b}}{b}$$

$$\frac{6}{\sqrt{6}}$$

$$= \frac{6}{\sqrt{6}} \cdot \frac{\sqrt{6}}{\sqrt{6}}$$

$$= \frac{6\sqrt{6}}{6}$$

$$= \sqrt{6}$$

Multiply the numerator and the denominator by the square root in the denominator and simplify

1. $\sqrt{\frac{9}{25}}$

2. $\sqrt{\frac{25}{36}}$

3. $\sqrt{\frac{25}{49}}$

4. $\sqrt{\frac{81}{100}}$

5. $\sqrt{\frac{3}{4}}$

6. $\sqrt{\frac{5}{9}}$

7. $\sqrt{\frac{64}{a^2}}$

8. $\sqrt{\frac{y^4}{16}}$

9. $\sqrt{\frac{1}{2}}$

10. $\sqrt{\frac{1}{3}}$

11. $\sqrt{\frac{4}{7}}$

12. $\sqrt{\frac{9}{11}}$

13. $\sqrt{\frac{1}{15}}$

14. $\sqrt{\frac{1}{14}}$

15. $\sqrt{\frac{4}{75}}$

16. $\sqrt{\frac{5}{12}}$

17. $\frac{2}{\sqrt{2}}$

18. $\frac{6}{\sqrt{3}}$

19. $\frac{10}{\sqrt{8}}$

20. $\frac{15}{\sqrt{27}}$

21. $\sqrt{\frac{x^2}{y}}$

22. $\sqrt{\frac{1}{a}}$

23. $\sqrt{\frac{1}{x}}$

24. $\sqrt{\frac{a^2}{b^3}}$

25. $\frac{\sqrt{a^5}}{\sqrt{a}}$

26. Find the width of a rectangle whose diagonal is 17 feet and length is 8 feet.

27. Find the diagonal of a rectangle whose length is 5 meters and whose width is 4 meters.

28. A 13-foot ladder is placed against the wall of a house. If the bottom of the ladder is 5 feet from the house, how far from the ground is the top of the ladder?

29. At an altitude of h feet above the sea or level ground, the distance d in miles that a person can see an object is found by using the equation

$$d = \sqrt{\frac{3h}{2}}$$

How far can someone see who is in a tower 96 feet above the ground?

Simplify the following expressions. Leave no radicals in the denominator. Assume that all variables represent positive real numbers. See examples 9-3 C and D.

Examples $\sqrt[3]{\frac{1}{27}}$

Solutions $= \frac{\sqrt[3]{1}}{\sqrt[3]{27}}$
 $= \frac{1}{3}$

$$\sqrt[3]{\frac{8a^5}{b^3}}$$

$$= \frac{\sqrt[3]{8a^5}}{\sqrt[3]{b^3}}$$

$$= \frac{\sqrt[3]{2^3 a^5}}{b}$$

$$= \frac{2a\sqrt[3]{a^2}}{b}$$

Rewrite as the cube root of the numerator over the cube root of the denominator and simplify

$$\frac{1}{\sqrt[5]{b^2}}$$

$$= \frac{1}{\sqrt[5]{b^2}} \cdot \frac{\sqrt[5]{b^3}}{\sqrt[5]{b^3}}$$

$$= \frac{\sqrt[5]{b^3}}{\sqrt[5]{b^5}}$$

$$= \frac{\sqrt[5]{b^3}}{b}$$

Multiply the numerator and the denominator by $\sqrt[5]{b^3}$

Perfect 5th root

$$\sqrt[5]{b^5} = b$$

30. $\sqrt[3]{\frac{8}{27}}$ 31. $\sqrt[3]{\frac{1}{8}}$ 32. $\sqrt[4]{\frac{16}{81}}$ 33. $\sqrt[3]{\frac{27}{125}}$ 34. $\sqrt[3]{\frac{a^2}{b^2}}$
35. $\sqrt[3]{\frac{3a^6}{b^3}}$ 36. $\sqrt[3]{\frac{x}{y^{12}}}$ 37. $\sqrt[5]{\frac{a^4}{b^{10}}}$ 38. $\sqrt[5]{\frac{32x^4}{y^5}}$ 39. $\sqrt[4]{\frac{a^4b^9}{c^{11}}}$
40. $\sqrt[4]{\frac{a^9b^{13}}{c^8}}$ 41. $\sqrt[5]{\frac{x^3y^2}{z^{15}}}$ 42. $\sqrt[3]{\frac{8}{9}}$ 43. $\sqrt[3]{\frac{4}{25}}$ 44. $\sqrt[3]{\frac{27}{16}}$
45. $\sqrt[4]{\frac{16}{125}}$ 46. $\sqrt[4]{\frac{3}{4}}$ 47. $\sqrt[3]{\frac{x^3}{y^2}}$ 48. $\sqrt[3]{\frac{x^6}{y}}$ 49. $\frac{ab}{\sqrt[3]{a^2}}$
50. $\frac{xy}{\sqrt[5]{y^3}}$ 51. $\sqrt[3]{\frac{a^3}{b^2c}}$ 52. $\sqrt[3]{\frac{8}{xy^2}}$ 53. $\sqrt[3]{\frac{a^2}{b^2c}}$ 54. $\frac{a}{\sqrt[5]{a^2b^4}}$
55. $\frac{ab}{\sqrt[3]{ab^2}}$ 56. $\frac{xy^2}{\sqrt[5]{x^4y}}$

57. If we wish to construct a sphere of specific volume, we can find the length of radius necessary by the formula $r = \sqrt[3]{\frac{3V}{4\pi}}$. Find the radius necessary for a sphere to have a volume of 113.04 cubic units. (Use 3.14 for π .)
58. Use exercise 57 to find r if $V = 904.32$ cubic units. (Use 3.14 for π .)

Review exercises

Combine in the following. See section 2-3.

1. $4x + 2x$ 2. $9y - 5y$ 3. $5ab + 3ab$ 4. $xy + 4xy$

Multiply the following. See section 3-2.

5. $(x + 3)(x - 3)$ 6. $(x + y)(x - y)$

9-4 ■ Sums and differences of radicals

Like radicals

We have learned that in addition and subtraction of algebraic expressions, we can only combine like terms. This same idea applies when we are dealing with radicals. **We can add or subtract only like radicals.** Like radicals are radicals having the same index and the same radicand. For example, $3\sqrt{5x}$ and $-2\sqrt{5x}$ are like radicals, but $5\sqrt{7x}$ and $7\sqrt{5x}$ are not, because the radicands are not the same.

Addition and subtraction involving square roots

Addition and subtraction of radicals follow the same procedure as addition and subtraction of algebraic expressions. That is, *once we have determined that we have like radicals, the operations of addition and subtraction are performed only with the numerical coefficients.*

Example 9-4 A

Perform the indicated operations and simplify. Assume that all variables represent nonnegative real numbers.

$$1. 5\sqrt{2} + 3\sqrt{2} = (5 + 3)\sqrt{2} = 8\sqrt{2}$$

Apply the distributive property

$$2. 12\sqrt{3} - \sqrt{3} = (12 - 1)\sqrt{3} = 11\sqrt{3}$$

$$3. 2\sqrt{a} + 3\sqrt{a} = 5\sqrt{a}$$

► **Quick check** Simplify. $3\sqrt{6} + 2\sqrt{6} - \sqrt{6}$

Consider the example

$$\sqrt{27} + 4\sqrt{3}$$

It appears that the indicated addition cannot be performed since we do not have like radicals. However we should have observed that the $\sqrt{27}$ can be simplified as

$$\sqrt{27} = \sqrt{9 \cdot 3} = 3\sqrt{3}$$

Our problem then becomes

$$\sqrt{27} + 4\sqrt{3} = 3\sqrt{3} + 4\sqrt{3} = 7\sqrt{3}$$

and we are able to add the like radicals. Therefore, *whenever we are working with radicals, we must be certain that all radicals are in simplest form.*

To combine like radicals

1. Perform any simplification within the terms.
2. Use the distributive property to combine terms that have like radicals.

Example 9-4 B

Perform the indicated operations. Assume that all variables represent nonnegative real numbers.

$$1. \sqrt{45} + \sqrt{20} = \sqrt{9 \cdot 5} + \sqrt{4 \cdot 5}$$

$$= 3\sqrt{5} + 2\sqrt{5}$$

$$= 5\sqrt{5}$$

Factor $45 = 9 \cdot 5$ and $20 = 4 \cdot 5$

$$\sqrt{4} = 2; \sqrt{9} = 3$$

Add coefficients

$$2. \sqrt{32} + 5\sqrt{8} = \sqrt{16 \cdot 2} + 5\sqrt{4 \cdot 2}$$

$$= 4\sqrt{2} + 5 \cdot 2\sqrt{2}$$

$$= 4\sqrt{2} + 10\sqrt{2}$$

$$= 14\sqrt{2}$$

Factor $32 = 16 \cdot 2$; $8 = 4 \cdot 2$

$$\sqrt{16} = 4 \text{ and } \sqrt{4} = 2$$

Multiply $5 \cdot 2 = 10$

Add coefficients

$$3. 3\sqrt{3a} - \sqrt{12a} + 5\sqrt{48a}$$

$$= 3\sqrt{3a} - \sqrt{4 \cdot 3a} + 5\sqrt{16 \cdot 3a}$$

$$= 3\sqrt{3a} - 2\sqrt{3a} + 5 \cdot 4\sqrt{3a}$$

$$= 3\sqrt{3a} - 2\sqrt{3a} + 20\sqrt{3a}$$

$$= 21\sqrt{3a}$$

Factor $12 = 4 \cdot 3$; $48 = 16 \cdot 3$

$$\sqrt{4} = 2; \sqrt{16} = 4$$

$$5 \cdot 4 = 20$$

Combine coefficients

► **Quick check** Simplify. $5\sqrt{2} + \sqrt{18}$

Addition and subtraction involving n th roots

Addition and subtraction of radicals other than square roots follow the same procedure as addition and subtraction of expressions containing square roots. That is, *once we have determined that we have like radicals, the operations of addition and subtraction are performed only with the numerical coefficients.*

Example 9-4 C

Perform the indicated operations and simplify. Assume that all variables represent nonnegative real numbers.

$$1. \sqrt[3]{5} + 6\sqrt[3]{5} = (1 + 6)\sqrt[3]{5} = 7\sqrt[3]{5}$$

Combine coefficients

$$\begin{aligned} 2. 4\sqrt[3]{81} - \sqrt[3]{24} &= 4\sqrt[3]{27 \cdot 3} - \sqrt[3]{8 \cdot 3} \\ &= 4 \cdot 3\sqrt[3]{3} - 2\sqrt[3]{3} \\ &= 12\sqrt[3]{3} - 2\sqrt[3]{3} \\ &= 10\sqrt[3]{3} \end{aligned}$$

Factor $81 = 27 \cdot 3$; $24 = 8 \cdot 3$ $\sqrt[3]{27} = 3$; $\sqrt[3]{8} = 2$ $4 \cdot 3 = 12$

Subtract coefficients

$$\begin{aligned} 3. \sqrt[3]{16x^2y} + \sqrt[3]{54x^2y} &= \sqrt[3]{8 \cdot 2x^2y} + \sqrt[3]{27 \cdot 2x^2y} \\ &= 2\sqrt[3]{2x^2y} + 3\sqrt[3]{2x^2y} \\ &= 5\sqrt[3]{2x^2y} \end{aligned}$$

Factor $16 = 8 \cdot 2$; $54 = 27 \cdot 2$ $\sqrt[3]{8} = 2$; $\sqrt[3]{27} = 3$

Add coefficients

Mastery points*Can you*

- Identify like radicals?
- Add and subtract like radicals?

Exercise 9-4

Perform the indicated operations and simplify. Assume that all variables represent nonnegative real numbers. See examples 9-4 A and B.

Examples $3\sqrt{6} + 2\sqrt{6} - \sqrt{6}$

Solutions $= (3 + 2 - 1)\sqrt{6}$
 $= 4\sqrt{6}$

Distributive property
Combine coefficients

$$5\sqrt{2} + \sqrt{18}$$

$$\begin{aligned} &= 5\sqrt{2} + \sqrt{9 \cdot 2} \\ &= 5\sqrt{2} + 3\sqrt{2} \\ &= (5 + 3)\sqrt{2} \\ &= 8\sqrt{2} \end{aligned}$$

Factor $18 = 9 \cdot 2$ $\sqrt{9} = 3$

Distributive property

Add coefficients

1. $5\sqrt{3} + 4\sqrt{3}$

4. $9\sqrt{6} - 6\sqrt{6}$

7. $\sqrt{7} + 5\sqrt{7} - 3\sqrt{7}$

10. $3\sqrt{x} + 4\sqrt{x}$

13. $5\sqrt{xy} + 2\sqrt{xy}$

16. $\sqrt{ab} + 2\sqrt{ab} + 3\sqrt{a}$

19. $\sqrt{8} + 5\sqrt{2}$

22. $2\sqrt{3} + 3\sqrt{12}$

25. $4\sqrt{2} - \sqrt{8} + \sqrt{50}$

2. $8\sqrt{7} - 2\sqrt{7}$

5. $2\sqrt{3} + 7\sqrt{3} - 3\sqrt{3}$

8. $2\sqrt{10} + 11\sqrt{10} - 9\sqrt{10}$

11. $5\sqrt{a} - 4\sqrt{a} + 7\sqrt{a}$

14. $3\sqrt{x} + 2\sqrt{y} - \sqrt{x}$

17. $5\sqrt{xy} - \sqrt{xy} + 3\sqrt{y}$

20. $\sqrt{12} + \sqrt{75}$

23. $5\sqrt{7} + 4\sqrt{63}$

26. $\sqrt{75} - 4\sqrt{3} + 2\sqrt{27}$

3. $6\sqrt{5} + 4\sqrt{5}$

6. $5\sqrt{5} - 4\sqrt{5} + 6\sqrt{5}$

9. $\sqrt{a} + 2\sqrt{a}$

12. $6\sqrt{y} - \sqrt{y} + 4\sqrt{y}$

15. $5\sqrt{a} + 2\sqrt{ab} + 3\sqrt{a}$

18. $\sqrt{20} + 3\sqrt{5}$

21. $\sqrt{48} - \sqrt{27}$

24. $5\sqrt{3} + \sqrt{27} - \sqrt{12}$

27. $\sqrt{12} + \sqrt{18} + \sqrt{50}$

28. $\sqrt{63} - \sqrt{28} + \sqrt{24}$

31. $3\sqrt{9x} - 5\sqrt{4x}$

34. $3\sqrt{48b} - 2\sqrt{12b} + \sqrt{3b}$

29. $\sqrt{50a} + \sqrt{8a}$

32. $2\sqrt{4x^2y} + 3\sqrt{25x^2y}$

35. $\sqrt{50a} + 3\sqrt{12a} - \sqrt{18a}$

30. $\sqrt{32a} - \sqrt{18a}$


33. $2\sqrt{8a} + 4\sqrt{50a} - 7\sqrt{2a}$

36. $4\sqrt{25x^2y} + 3\sqrt{81x^2y} - 2\sqrt{2y}$


37. We can find the height, h , of the given figure by finding b from the formula $b = \sqrt{c^2 - s^2}$.

If $c = 10$ units and $s = 6$ units, find h .

38. Use exercise 37 to find the height of the figure if $c = 13$ feet and $s = 5$ feet.



39. The figure is made up of 9 equal squares in which each square has an area of 7.29 square units. What are the dimensions of the figure?



Perform the indicated operations and simplify. Assume that all variables represent nonnegative real numbers. See example 9-4 C.

40. $3\sqrt[3]{4} + 5\sqrt[3]{4}$

41. $7\sqrt[5]{2} - 4\sqrt[5]{2} + 3\sqrt[5]{2}$

42. $9\sqrt[4]{3} + 6\sqrt[4]{3} + 2\sqrt[4]{3}$

43. $\sqrt[3]{16} + \sqrt[3]{54}$

44. $\sqrt[3]{24} - \sqrt[3]{81}$

45. $\sqrt[3]{81} + 2\sqrt[3]{250}$

46. $\sqrt[3]{8a^2} + \sqrt[3]{27a^2}$

47. $\sqrt[4]{16x^3} + \sqrt[4]{81x^3}$

48. $\sqrt[4]{625a} - \sqrt[4]{81a}$

49. $\sqrt[3]{64x^2y} - \sqrt[3]{27x^2y}$

50. $\sqrt[3]{x^6y} + 2x^2\sqrt[3]{y}$

51. $3a\sqrt[3]{b^2} - \sqrt[3]{a^3b^2}$

52. $\sqrt[3]{16a^2b} + \sqrt[3]{54a^2b}$

Review exercises

Multiply the following expressions. See section 3-2.

1. $3x(2x - y)$

2. $2a^2(a^2 - b^2)$

3. $(x - 1)(x - 1)$

4. $(y - 1)(y + 1)$

5. $(2x + 1)(2x - 1)$

6. $(x + 3y)(x + 2y)$

7. $(x - y)^2$

8. $(a + 2b)^2$

9-5 Further operations with radicals

Multiplication of radical expressions

In section 9-2, we learned the procedure for multiplying two radicals. We now combine those ideas along with the *distributive property*, $a(b + c) = ab + ac$, to perform multiplication of radical expressions containing more than one term.

Example 9-5 A

Perform the indicated operations and simplify. Assume that all variables represent nonnegative real numbers.

$$\begin{aligned} 1. \sqrt{3}(3 + \sqrt{3}) &= \sqrt{3} \cdot 3 + \sqrt{3}\sqrt{3} \\ &= 3\sqrt{3} + \sqrt{9} \\ &= 3\sqrt{3} + 3 \end{aligned}$$

Distributive property
 $\sqrt{3}\sqrt{3} = \sqrt{9}$
 $\sqrt{9} = 3$

$$\begin{aligned} 2. \sqrt{3}(\sqrt{6} + \sqrt{21}) &= \sqrt{3}\sqrt{6} + \sqrt{3}\sqrt{21} \\ &= \sqrt{18} + \sqrt{63} \\ &= \sqrt{9 \cdot 2} + \sqrt{9 \cdot 7} \\ &= 3\sqrt{2} + 3\sqrt{7} \end{aligned}$$

Distributive property
 Multiply radicands
 Factor $18 = 9 \cdot 2$;
 $63 = 9 \cdot 7$
 Simplify radicals

$$3. (\sqrt{2} + \sqrt{3})(\sqrt{2} + 5\sqrt{3})$$

Note In this example, we are multiplying two binomials. Therefore, as we did in chapter 3, we will *multiply each term in the first parentheses by each term in the second parentheses*.

$$\begin{aligned} &= \sqrt{2}\sqrt{2} + \sqrt{2} \cdot 5\sqrt{3} + \sqrt{3}\sqrt{2} + \sqrt{3} \cdot 5\sqrt{3} \\ &= \sqrt{4} + 5\sqrt{6} + \sqrt{6} + 5 \cdot \sqrt{9} \\ &= 2 + 5\sqrt{6} + \sqrt{6} + 5 \cdot 3 \\ &= 2 + 5\sqrt{6} + \sqrt{6} + 15 \\ &= 17 + 6\sqrt{6} \end{aligned}$$

Distributive property
 Multiply radicands
 $\sqrt{4} = 2$, $\sqrt{9} = 3$

$$\begin{aligned} 4. (3 - \sqrt{2})(3 + \sqrt{2}) &= 9 + 3\sqrt{2} - 3\sqrt{2} - \sqrt{4} \\ &= 9 + 3\sqrt{2} - 3\sqrt{2} - 2 \\ &= 9 - 2 \\ &= 7 \end{aligned}$$

Combine like terms
 Distributive property
 Simplify radicals
 Combine like terms
 Subtract

We observe that when we simplified, there were no longer any radicals in the answer.

$$\begin{aligned} 5. (\sqrt{a} - \sqrt{b})(\sqrt{a} + \sqrt{b}) &= \sqrt{a}\sqrt{a} + \sqrt{a}\sqrt{b} - \sqrt{b}\sqrt{a} - \sqrt{b}\sqrt{b} \\ &= \sqrt{a^2} + \sqrt{ab} - \sqrt{ab} - \sqrt{b^2} \\ &= a + \sqrt{ab} - \sqrt{ab} - b \\ &= a - b \end{aligned}$$

Distributive property
 Multiply radicands
 $\sqrt{a^2} = a$ and $\sqrt{b^2} = b$
 Combine like terms

$$\begin{aligned} 6. (\sqrt{3} + 2\sqrt{2})^2 &= (\sqrt{3} + 2\sqrt{2})(\sqrt{3} + 2\sqrt{2}) \\ &= \sqrt{3}\sqrt{3} + \sqrt{3} \cdot 2\sqrt{2} + \sqrt{3} \cdot 2\sqrt{2} + 2\sqrt{2} \cdot 2\sqrt{2} \\ &= \sqrt{9} + 2\sqrt{6} + 2\sqrt{6} + 4\sqrt{4} \\ &= 3 + 2\sqrt{6} + 2\sqrt{6} + 4 \cdot 2 \\ &= 3 + 2\sqrt{6} + 2\sqrt{6} + 8 \\ &= 11 + 4\sqrt{6} \end{aligned}$$

Distributive property
 Multiply radicands
 Simplify radicals
 Multiply
 Combine like terms

► **Quick check** Simplify. $2(\sqrt{3} + \sqrt{5})$; $\sqrt{2}(\sqrt{14} + \sqrt{6})$; and $(\sqrt{2} + \sqrt{3})(\sqrt{2} - 2\sqrt{3})$

Conjugate factors

The type of factors that we are multiplying in examples 4 and 5 are called **conjugate factors**. The conjugate is used to rationalize the denominator of a fraction when the denominator contains two terms where one or both terms contain a square root. The idea of conjugate factors is derived from the factorization of the difference of two squares. When multiplying conjugate factors, we can simply write our answer as the square of the second term subtracted from the square of the first term.

In examples 4 and 5, we could have performed the multiplication as follows:

$$4. (3 - \sqrt{2})(3 + \sqrt{2}) = (3)^2 - (\sqrt{2})^2 = 9 - 2 = 7;$$

$$5. (\sqrt{a} - \sqrt{b})(\sqrt{a} + \sqrt{b}) = (\sqrt{a})^2 - (\sqrt{b})^2 = a - b.$$

To determine what the conjugate of a given factor is, we write the original factor and change the sign of the second term.

■ Example 9-5 B

Form the conjugates of the given expressions.

- | | |
|---------------------------|---|
| 1. $\sqrt{7} + 2$ | The conjugate is $\sqrt{7} - 2$. |
| 2. $\sqrt{11} - \sqrt{6}$ | The conjugate is $\sqrt{11} + \sqrt{6}$. |
| 3. $-5 - 2\sqrt{3}$ | The conjugate is $-5 + 2\sqrt{3}$. |
| 4. $\sqrt{a} + \sqrt{b}$ | The conjugate is $\sqrt{a} - \sqrt{b}$. |

► **Quick check** Form the conjugate. $6 - 3\sqrt{2}$

Rationalizing the denominator

If we wish to rationalize the denominator of the fraction

$$\frac{1}{3 - \sqrt{2}}$$

we recall from example 9-5 A-4 that when we multiplied $3 - \sqrt{2}$ by $3 + \sqrt{2}$, there were no radicals left in our product. This result is precisely what we want to occur in our denominator. Therefore, to rationalize this fraction, we apply the fundamental principle of fractions and multiply the numerator and the denominator by $3 + \sqrt{2}$, the conjugate of the denominator.

$$\begin{aligned} \frac{1}{3 - \sqrt{2}} &= \frac{1}{3 - \sqrt{2}} \cdot \frac{3 + \sqrt{2}}{3 + \sqrt{2}} && 3 + \sqrt{2} \text{ is the conjugate of the denominator} \\ &= \frac{1(3 + \sqrt{2})}{(3)^2 - (\sqrt{2})^2} && (\text{first term})^2 - (\text{second term})^2 \\ &= \frac{3 + \sqrt{2}}{9 - 2} && \text{No radicals remain in the denominator} \\ &= \frac{3 + \sqrt{2}}{7} && \text{Denominator is rationalized} \end{aligned}$$

Example 9-5 C

Rationalize the denominators.

$$\begin{aligned}
 1. \quad \frac{2}{\sqrt{7} + 2} &= \frac{2}{\sqrt{7} + 2} \cdot \frac{\sqrt{7} - 2}{\sqrt{7} - 2} \\
 &= \frac{2(\sqrt{7} - 2)}{(\sqrt{7})^2 - (2)^2} \\
 &= \frac{2\sqrt{7} - 4}{7 - 4} \\
 &= \frac{2\sqrt{7} - 4}{3}
 \end{aligned}$$

Multiply by the conjugate of the denominator

$$(x + y)(x - y) = x^2 - y^2$$

Simplify in numerator and denominator

Subtract in denominator

$$\begin{aligned}
 2. \quad \frac{5}{\sqrt{11} - \sqrt{6}} &= \frac{5}{\sqrt{11} - \sqrt{6}} \cdot \frac{\sqrt{11} + \sqrt{6}}{\sqrt{11} + \sqrt{6}} \\
 &= \frac{5(\sqrt{11} + \sqrt{6})}{(\sqrt{11})^2 - (\sqrt{6})^2} \\
 &= \frac{5(\sqrt{11} + \sqrt{6})}{11 - 6} \\
 &= \frac{5(\sqrt{11} + \sqrt{6})}{5} \\
 &= \sqrt{11} + \sqrt{6}
 \end{aligned}$$

Multiply by the conjugate of the denominator

$$(x + y)(x - y) = x^2 - y^2$$

Simplify radicals

Subtract in denominator

Reduce (by 5) to lowest terms

$$\begin{aligned}
 3. \quad \frac{\sqrt{3}}{5 - 2\sqrt{3}} &= \frac{\sqrt{3}}{5 - 2\sqrt{3}} \cdot \frac{5 + 2\sqrt{3}}{5 + 2\sqrt{3}} \\
 &= \frac{\sqrt{3}(5 + 2\sqrt{3})}{(5)^2 - (2\sqrt{3})^2} \\
 &= \frac{5\sqrt{3} + 2\sqrt{9}}{25 - 2^2(\sqrt{3})^2} \\
 &= \frac{5\sqrt{3} + 2 \cdot 3}{25 - 4 \cdot 3} \\
 &= \frac{5\sqrt{3} + 6}{25 - 12} \\
 &= \frac{5\sqrt{3} + 6}{13}
 \end{aligned}$$

Multiply by the conjugate of the denominator

$$(x + y)(x - y) = x^2 - y^2$$

Simplify radicals

Simplify radicals

Perform operations

Subtract in denominator

► **Quick check** Rationalize the denominators. $\frac{3}{7 + \sqrt{2}}$ and $\frac{2}{\sqrt{11} - 3}$

Mastery points*Can you*

- Multiply radical expressions containing more than one term?
- Form conjugate factors?
- Multiply conjugate factors?
- Rationalize a denominator that has two terms in which one or both terms contain a square root?

Exercise 9-5

Perform the indicated operations and simplify. Assume that all variables represent positive real numbers. See example 9-5 A.

Examples $2(\sqrt{3} + \sqrt{5})$

Solutions $= 2\sqrt{3} + 2\sqrt{5}$

Distributive property

$\sqrt{2}(\sqrt{14} + \sqrt{6})$

$$\begin{aligned} &= \sqrt{2}\sqrt{14} + \sqrt{2}\sqrt{6} \\ &= \sqrt{28} + \sqrt{12} \\ &= \sqrt{4 \cdot 7} + \sqrt{4 \cdot 3} \\ &= 2\sqrt{7} + 2\sqrt{3} \end{aligned}$$

Distributive property

Product property

Factor $28 = 4 \cdot 7$; $12 = 4 \cdot 3$

$\sqrt{4} = 2$

Example $(\sqrt{2} + \sqrt{3})(\sqrt{2} - 2\sqrt{3})$

Solution
$$\begin{aligned} &= \sqrt{2}\sqrt{2} - \sqrt{2} \cdot 2\sqrt{3} + \sqrt{3}\sqrt{2} - \sqrt{3} \cdot 2\sqrt{3} \\ &= 2 - 2\sqrt{6} + \sqrt{6} - 2 \cdot 3 \\ &= 2 - \sqrt{6} - 6 \\ &= -4 - \sqrt{6} \end{aligned}$$

Distributive property

$\sqrt{2}\sqrt{2} = 2$; $\sqrt{3}\sqrt{3} = 3$

Combine like radicals

$2 - 6 = -4$

1. $3(\sqrt{2} + \sqrt{3})$

4. $\sqrt{5}(\sqrt{7} - \sqrt{3})$

7. $\sqrt{5}(\sqrt{15} - \sqrt{10})$

10. $\sqrt{a}(\sqrt{ab} + \sqrt{b})$

13. $(3 + \sqrt{2})(4 + \sqrt{2})$

16. $(7 + 2\sqrt{y})(6 + 5\sqrt{y})$

19. $(2 + \sqrt{6})(2 - \sqrt{6})$

22. $(3 - \sqrt{7})^2$

25. $(\sqrt{x} - \sqrt{y})(\sqrt{x} + \sqrt{y})$

28. $(a\sqrt{b} + c)(a\sqrt{b} - c)$

2. $5(2\sqrt{6} + \sqrt{2})$

5. $3\sqrt{2}(2\sqrt{3} - \sqrt{11})$

8. $\sqrt{14}(\sqrt{21} + \sqrt{10})$

11. $\sqrt{a}(3\sqrt{a} + \sqrt{b})$

14. $(5 - \sqrt{5})(5 - \sqrt{5})$

17. $(\sqrt{3} + \sqrt{2})(\sqrt{3} - \sqrt{2})$

20. $(5 - \sqrt{3})(5 + \sqrt{3})$

23. $(\sqrt{x} + \sqrt{y})^2$

26. $(2\sqrt{a} - \sqrt{b})(2\sqrt{a} + \sqrt{b})$

29. $(2\sqrt{x} + y)^2$

3. $\sqrt{2}(\sqrt{3} + \sqrt{7})$

6. $\sqrt{6}(\sqrt{2} + \sqrt{3})$

9. $2\sqrt{7}(\sqrt{35} - 3\sqrt{14})$

12. $(5 + \sqrt{3})(4 - \sqrt{3})$

15. $(3 - 4\sqrt{a})(4 - 3\sqrt{a})$

18. $(\sqrt{7} + \sqrt{5})(\sqrt{7} - \sqrt{5})$

21. $(2 + \sqrt{5})^2$

24. $(\sqrt{a} - \sqrt{b})^2$

27. $(x\sqrt{y} + \sqrt{z})(x\sqrt{y} - \sqrt{z})$

30. $(3\sqrt{a} + \sqrt{b})^2$

Form the conjugate of the given expressions. See example 9-5 B.

Example $6 - 3\sqrt{2}$

Solution $6 + 3\sqrt{2}$

First term remains the same, change the sign of the second term

31. $11 - \sqrt{3}$

32. $-5\sqrt{7} - \sqrt{2}$

33. $\sqrt{a} + 3\sqrt{b}$

34. $a\sqrt{b} - \sqrt{c}$

Rationalize the denominators. Assume that all variables represent positive real numbers and that no denominator is equal to zero. See example 9-5 C.

Love The Taste. Taste The Love.

At Culver's® we can't think of anything better than serving up our creamy frozen custard and delicious classics cooked fresh the minute you order them. Which is why when we bring them to your table, they're always accompanied by a warm smile and a friendly offer to see if there's anything else we can get for you. So come on into your neighborhood Culver's and see for yourself. You might just be in love by the time you leave.



Simplify the following expressions, leaving all denominators rationalized. Assume that all variables represent positive real numbers and that no denominator is equal to zero. See example 9-5 C.

Examples	$\frac{3}{7 + \sqrt{2}}$		$\frac{2}{\sqrt{11} - 3}$	
Solutions	$= \frac{3}{7 + \sqrt{2}} \cdot \frac{7 - \sqrt{2}}{7 - \sqrt{2}}$	Multiply by the conjugate	$= \frac{2}{\sqrt{11} - 3} \cdot \frac{\sqrt{11} + 3}{\sqrt{11} + 3}$	Multiply by the conjugate
	$= \frac{3(7 - \sqrt{2})}{(7)^2 - (\sqrt{2})^2}$	$(x + y)(x - y) = x^2 - y^2$	$= \frac{2(\sqrt{11} + 3)}{(\sqrt{11})^2 - (3)^2}$	$(x + y)(x - y) = x^2 - y^2$
	$= \frac{3(7 - \sqrt{2})}{49 - 2}$	Simplify denominator	$= \frac{2(\sqrt{11} + 3)}{11 - 9}$	Simplify denominator
	$= \frac{3(7 - \sqrt{2})}{47}$	Simplify denominator	$= \frac{2(\sqrt{11} + 3)}{2}$	Simplify denominator
	$= \frac{21 - 3\sqrt{2}}{47}$	Multiply in numerator	$= \sqrt{11} + 3$	Reduce fraction

35. $\frac{1}{\sqrt{2} + 3}$

36. $\frac{1}{\sqrt{3} - 2}$

37. $\frac{7}{2 + \sqrt{7}}$

38. $\frac{6}{3 - \sqrt{6}}$

39. $\frac{3}{\sqrt{6} - \sqrt{3}}$

40. $\frac{1}{\sqrt{a} + b}$

41. $\frac{3}{2\sqrt{3} - \sqrt{5}}$

42. $\frac{4}{2\sqrt{3} - \sqrt{6}}$

43. $\frac{1 + \sqrt{5}}{1 - \sqrt{5}}$

44. $\frac{\sqrt{3} - \sqrt{7}}{\sqrt{3} + \sqrt{7}}$

45. $\frac{\sqrt{a} + b}{\sqrt{a} - b}$

Review exercises

Perform the indicated operations and leave your answer with only positive exponents. Assume that no variable is equal to zero. See sections 3-1 and 3-3.

1. $(2^2)^3$

2. $2^2 \cdot 2^3$

3. $3^{-2} \cdot 3^4$

4. $\frac{3^2}{3^5}$

5. $(x^{-2})^3$

6. $(2a^2b)^3$

7. $\frac{x^2y^5}{x^3y^2}$

8. $x^{-4} \cdot x^{-5}$

9-6 ■ Fractional exponents

Fractional exponents

In this section, we are going to develop the idea of a fraction used as an exponent. Consider the example

$$(a^{1/2})^2 = a^{1/2 \cdot 2} = a^1$$

When we raise a power to a power, we multiply the exponents. In the previous section, we observed that if a represented only nonnegative real numbers, then

$$(\sqrt{a})^2 = a$$

Therefore, for our properties of exponents and our procedures for radicals to be consistent, the following statement must be true:

$$a^{1/2} = \sqrt{a}$$

We generalize as follows:

Definition of $a^{1/n}$

$$a^{1/n} = \sqrt[n]{a}$$

where n is a natural number greater than 1. Whenever n is even, a represents only nonnegative real numbers.

Concept

The expression $a^{1/n}$ represents the principal n th root of a .

Example 9-6 A

Rewrite the following in radical notation and simplify where possible.

1. $5^{1/2} = \sqrt{5}$

2. $(64)^{1/2} = \sqrt{64} = 8$

3. $a^{1/3} = \sqrt[3]{a}$

4. $(-8)^{1/3} = \sqrt[3]{(-8)} = -2$

Once we become acquainted with fractional exponents, the process of changing the fractional exponent to radical form for simplification will become unnecessary.

Consider the following problem:

$$(\sqrt[3]{a})^2 = (a^{1/3})^2 = a^{2/3}$$

We observe that *when a number is raised to a fractional exponent, the numerator of the fractional exponent indicates the power to which the base is to be raised, and the denominator indicates the root to be taken.*

Definition of $a^{m/n}$

If a is any real number, m is any integer, and n is any positive integer, then if $\frac{m}{n}$ is reduced to lowest terms,

$$a^{m/n} = (\sqrt[n]{a})^m = \sqrt[n]{a^m}$$

provided that $\sqrt[n]{a}$ is a real number.

Concept

The expression $a^{m/n}$ represents the principal n th root of a raised to the m th power.

We calculate $a^{m/n}$ by first finding the principal n th root of a and then raising the resulting number to the m th power.

Example 9-6 B

Simplify. Assume that all variables represent nonnegative real numbers.

Numerator is the power Denominator is the index

$$1. (16)^{3/4} = (\sqrt[4]{16})^3$$

$$= 2^3$$

$$= 8$$

4th root of 16 is 2
Standard form

$$2. (32)^{2/5} = (\sqrt[5]{32})^2 = 2^2 = 4$$

$$3. (-27)^{2/3} = (\sqrt[3]{-27})^2 = (-3)^2 = 9$$

$$4. (x^6)^{1/3} = \sqrt[3]{x^6} = x^2$$

► **Quick check** Simplify. $(49)^{1/2}$ and $(8)^{2/3}$

Operations with fractional exponents

We can extend the properties and definitions involving integer exponents to expressions that involve fractional exponents.

Example 9-6 C

Perform the indicated operations and simplify. Assume that all variables represent positive real numbers.

$$1. 5^{1/2} \cdot 5^{1/3} = 5^{1/2 + 1/3}$$

Multiply like bases, add exponents

$$= 5^{3/6 + 2/6}$$

Least common denominator is 6

$$= 5^{5/6}$$

Add numerators

$$2. \frac{2^{1/2}}{2^{1/4}} = 2^{1/2 - 1/4} = 2^{2/4 - 1/4} = 2^{1/4}$$

$$3. (-27)^{-2/3} = \frac{1}{(-27)^{2/3}} = \frac{1}{(\sqrt[3]{-27})^2} = \frac{1}{(-3)^2} = \frac{1}{9}$$

$$4. (2^3 x^9 y^{15})^{1/3} = (2^3)^{1/3} (x^9)^{1/3} (y^{15})^{1/3} = 2^3 \cdot 1/3 \cdot x^9 \cdot 1/3 \cdot y^{15} \cdot 1/3 = 2^1 x^3 y^5$$

$$= 2x^3y^5$$

$$5. x^{2/3} \cdot x^{3/4} = x^{2/3 + 3/4} = x^{8/12 + 9/12} = x^{17/12}$$

$$6. (y^{1/2})^{4/3} = y^{1/2 \cdot 4/3} = y^{2/3}$$

$$7. \frac{z^{1/2}}{z^{2/3}} = z^{1/2 - 2/3} = z^{3/6 - 4/6} = z^{-1/6} = \frac{1}{z^{1/6}}$$

Note In example 7, $z^{1/6}$ in the denominator is simply another form of $\sqrt[6]{z}$. Therefore, if we want our answer in a rationalized form, we would proceed as follows:

$$\frac{1}{z^{1/6}} = \frac{1}{\sqrt[6]{z}} \cdot \frac{\sqrt[6]{z^5}}{\sqrt[6]{z^5}} = \frac{\sqrt[6]{z^5}}{z} = \frac{z^{5/6}}{z}$$

► **Quick check** Simplify. $y^{2/3} \cdot y^{3/2}$; $\frac{a^{3/4}b^{5/3}}{a^{1/4}b}$; and $(36)^{-1/2}$

Mastery points**Can you**

- Express fractional exponents in radical form?
- Express radicals in fractional exponent form?
- Apply the properties and definitions involving integer exponents to fractional exponents?

Exercise 9-6

Simplify the given expressions. See examples 9-6 A and B.

Examples $(49)^{1/2}$ **Solutions** $= \sqrt{49}$
 $= 7$

Square root

$$\begin{aligned} 8^{2/3} &= (\sqrt[3]{8})^2 \\ &= (2)^2 \\ &= 4 \end{aligned}$$

Numerator is the power
Denominator is the index
Cube root of 8 is 2
Standard form

 $(36)^{-1/2}$

$$\begin{aligned} &= \frac{1}{(36)^{1/2}} \\ &= \frac{1}{\sqrt{36}} \\ &= \frac{1}{6} \end{aligned}$$

 $a^{-n} = \frac{1}{a^n}$

Square root

Simplify

1. $(36)^{1/2}$

2. $(25)^{1/2}$

3. $(a^6)^{1/3}$

4. $(b^{12})^{1/3}$

5. $(8)^{1/3}$

6. $(32)^{1/5}$

7. $(-27)^{1/3}$

8. $(-8)^{1/3}$

9. $(27)^{2/3}$

10. $(16)^{3/4}$

11. $(9)^{3/2}$

12. $(16)^{3/2}$

Perform the indicated operations and simplify. Assume that all variables represent positive real numbers. See example 9-6 C.

Examples $y^{2/3} \cdot y^{3/2}$

Solutions $= y^{2/3 + 3/2}$
 $= y^{4/6 + 9/6}$
 $= y^{13/6}$

Multiply like bases,
add exponents
Least common denominator is 6
Add numerators

$$\frac{a^{3/4}b^{5/3}}{a^{1/4}b}$$

$$\begin{aligned} &= a^{3/4 - 1/4}b^{5/3 - 1} \\ &= a^{2/4}b^{5/3 - 3/3} \\ &= a^{1/2}b^{2/3} \end{aligned}$$

Divide like bases,
subtract exponents
Common denominators
Subtract numerators

13. $(25)^{-1/2}$

14. $(9)^{-1/2}$

15. $(16)^{-3/4}$

16. $(27)^{-2/3}$

17. $(-8)^{-1/3}$

18. $(-27)^{-1/3}$

19. $2^{1/2} \cdot 2^{3/2}$

20. $3^{1/3} \cdot 3^{2/3}$

21. $2^{1/3} \cdot 2^{1/2}$

22. $5^{1/5} \cdot 5^{1/2}$

23. $x^{1/4} \cdot x^{3/4}$

24. $b^{1/3} \cdot b^{2/3}$

25. $c^{1/2} \cdot c^{1/4}$

26. $x^{1/4} \cdot x^{1/3}$

27. $\frac{2^{3/2}}{2^{1/2}}$

28. $\frac{3^{4/3}}{3^{1/3}}$

29. $\frac{2^{1/2}}{2^{1/3}}$

30. $\frac{7^{3/4}}{7^{2/3}}$

31. $\frac{a^{4/5}}{a^{1/5}}$

32. $\frac{x^{3/4}}{x^{1/2}}$

33. $\frac{y^{2/3}}{y^{1/2}}$

34. $\frac{x^{5/6}}{x^{2/3}}$

35. $(a^{2/3})^{1/2}$

36. $(c^{1/2})^{1/2}$

37. $(x^{1/2})^{4/3}$

38. $(y^{2/3})^{3/4}$

39. $(c^{-1/4})^{-2/3}$

40. $(y^{-1/2})^{-2/5}$

41. $(a^{-2/3})^{-1/2}$

42. $(b^{-1/2})^{-1/2}$

43. $(x^{1/4})^{-2/3}$

44. $(x^{1/3})^{-3/4}$

45. $(y^{-3/4})^{1/3}$

46. $(c^{-2/5})^{1/2}$

47. $(16a^4)^{3/4}$

48. $(x^3y^{12})^{1/3}$

49. $(8a^6b^3)^{2/3}$

50. $(27x^3y^{12})^{2/3}$

51. $\frac{b^{3/4}c^{1/2}}{b^{1/4}c^{1/4}}$

52. $\frac{xy^{3/4}}{x^{1/2}y^{1/4}}$

53. $\frac{ab}{a^{1/2}b^{1/3}}$

54. $\frac{xy^{3/4}}{x^{2/5}y^{1/2}}$

Solve the following word problems.

55. Find the number whose principal fourth root is 3.

56. Find the number whose principal cube root is -2 .

57. The formula for approximating the velocity V in miles per hour of a car based on the length of its skid marks S (in feet) on dry pavement is given by $V = 4.9 S^{1/2}$

If the skid marks are 100 feet long, what was the approximate velocity?

58. The formula for approximating the velocity V in miles per hour of a car based on the length of its skid marks S (in feet) on wet pavement is given by $V = 3.46 S^{1/2}$

If the skid marks are 81 feet long, what was the approximate velocity of the car?

59. At an altitude of h feet above the sea or level ground, the distance d in miles that a person can see an object is found by using the equation

$$d = 1.2 h^{1/2}$$

How far can someone see who is in a tower 400 feet above the ground?

60. How can you find the principal fourth root of a number on a calculator using only the square root key?

61. How can you find the principal eighth root of a number on a calculator using only the square root key?

Review exercises

Perform the indicated operations. Assume that all radicands are nonnegative. See sections 9-1 and 9-2.

1. $(\sqrt{7})^2$

2. $(\sqrt{x})^2$

3. $(\sqrt{x+1})^2$

Perform the indicated operations. See section 3-2.

4. $(x+1)^2$

5. $(x-2)^2$

Find the solution set for the following equations. See section 4-7.

6. $x+6=x^2$

7. $x+2=x^2-9x+18$

8. $x+1=x^2+2x+1$

9-7 ■ Equations involving radicals

Radical equations

An equation in which the unknown quantity appears under a radical symbol is called a **radical equation**. Examples of radical equations are

$$\sqrt{x} = 5; \quad \sqrt{x+2} = 7; \quad 4 + \sqrt{x+2} = x$$

In this section, we will consider radical equations containing only square roots.

Solving radical equations involves squaring both members of an equation to eliminate the square roots. When we square both members of an equation, we use the squaring property of equality.

Squaring property of equality

If P and Q are algebraic expressions and if

$$P = Q$$

then all solutions of $P = Q$ are also solutions of the equation

$$P^2 = Q^2$$

Concept

If each member of an equation is squared, the solution(s) of the original equation are solution(s) of the resulting equation.

Extraneous solutions

This property implies that there *may be* solutions of the equation $P^2 = Q^2$ that are not solutions of the original equation $P = Q$. If such solutions exist, they are called **extraneous solutions** (roots). Thus, all possible solutions must be checked in the original equation.

To solve a radical equation

1. Rewrite the equation (if necessary) so that a radical is by itself in one member of the equation.
2. Square each member (side) of the equation and combine like terms.
3. Repeat steps 1 and 2 if a radical remains in the equation.
4. Solve the resulting equation.
5. Check all possible solutions in the original equation.

Example 9-7 A

Find the solution set.

$$\begin{aligned} 1. \quad & \sqrt{x} = 5 \\ & (\sqrt{x})^2 = (5)^2 \\ & x = 25 \end{aligned}$$

Check:

$$\begin{aligned} & \sqrt{x} = 5 \\ & \sqrt{(25)} = 5 \\ & 5 = 5 \end{aligned}$$

The solution set is $\{25\}$.

$$\begin{aligned} 2. \quad & \sqrt{x+2} = 7 \\ & (\sqrt{x+2})^2 = (7)^2 \\ & x+2 = 49 \\ & x = 47 \end{aligned}$$

Check:

$$\begin{aligned} & \sqrt{x+2} = 7 \\ & \sqrt{(47)+2} = 7 \\ & \sqrt{49} = 7 \\ & 7 = 7 \end{aligned}$$

The solution set is $\{47\}$.

Square both members

Squaring a square root gives the radicand

Original equation

Substitute into original equation

True

Square both members

Squaring a square root gives the radicand

Subtract 2 from both members

Original equation

Substitute into original equation

Simplify the radicand

True

$$3. \quad 4 + \sqrt{x+2} = x$$

$$\sqrt{x+2} = x - 4$$

$$(\sqrt{x+2})^2 = (x-4)^2$$

$$x+2 = (x-4)(x-4)$$

$$x+2 = x^2 - 4x - 4x + 16$$

$$x+2 = x^2 - 8x + 16$$

$$0 = x^2 - 9x + 14$$

$$0 = (x-7)(x-2)$$

$$x-7 = 0 \text{ or } x-2 = 0$$

$$x = 7 \text{ or } x = 2$$

Isolate the radical by subtracting 4

Square both members

Simplify

Multiply

Subtract x and 2

Solve the resulting quadratic equation

Factor

Set each factor equal to 0 and solve

Check for extraneous roots

Check:

for 7

$$4 + \sqrt{x+2} = x$$

$$4 + \sqrt{(7)+2} = (7)$$

$$4 + \sqrt{9} = 7$$

$$4 + 3 = 7$$

$$7 = 7 \quad \text{True}$$

for 2

$$4 + \sqrt{x+2} = x$$

$$4 + \sqrt{(2)+2} = (2)$$

$$4 + \sqrt{4} = 2$$

$$4 + 2 = 2$$

$$6 = 2$$

Original equation

Substitute

Simplify

Simplify

False

Therefore, 7 is the only solution to the equation. 2 is an *extraneous root* and the solution set is $\{7\}$.

$$4. \quad \sqrt{x+1} = x+1$$

$$(\sqrt{x+1})^2 = (x+1)^2$$

$$x+1 = (x+1)(x+1)$$

$$x+1 = x^2 + x + x + 1$$

$$x+1 = x^2 + 2x + 1$$

$$0 = x^2 + x$$

$$0 = x(x+1)$$

$$x = 0 \text{ or } x+1 = 0$$

Therefore, $x = 0$ or $x = -1$.

Square both members

Simplify

Multiply

Combine like terms

Solve the resulting quadratic equation

Factor

Set each factor equal to 0 and solve

Check:

for 0

$$\sqrt{x+1} = x+1$$

$$\sqrt{(0)+1} = (0)+1$$

$$\sqrt{1} = 1$$

$$1 = 1 \quad \text{True}$$

for -1

$$\sqrt{x+1} = x+1$$

$$\sqrt{(-1)+1} = (-1)+1$$

$$\sqrt{0} = 0$$

$$0 = 0$$

Original equation

Substitute

Simplify

True

Therefore, 0 and -1 are solutions of the equation. The solution set is $\{0, -1\}$.

$$5. \quad \sqrt{3x+4} = \sqrt{x+14}$$

$$(\sqrt{3x+4})^2 = (\sqrt{x+14})^2$$

$$3x+4 = x+14$$

$$2x+4 = 14$$

$$2x = 10$$

$$x = 5$$

Square both members

Solve for x Subtract x from both members

Subtract 4 from both members

Divide both members by 2

Check:

$$\sqrt{3x+4} = \sqrt{x+14}$$

$$\sqrt{3(5)+4} = \sqrt{(5)+14}$$

$$\sqrt{15+4} = \sqrt{19}$$

$$\sqrt{19} = \sqrt{19}$$

The solution set is $\{5\}$.

Original equation

Substitute

Simplify

True

Note There are two square roots in this problem, but we can eliminate both radical symbols by squaring both members.

$$\begin{aligned}
 6. \quad & \sqrt{x+3} = -6 \\
 & (\sqrt{x+3})^2 = (-6)^2 && \text{Square both members} \\
 & x+3 = 36 && \text{Solve for } x \\
 & x = 33 && \text{Subtract 3 from both members} \\
 \text{Check:} & \\
 & \sqrt{x+3} = -6 && \text{Original equation} \\
 & \sqrt{(33)+3} = -6 && \text{Substitute} \\
 & \sqrt{36} = -6 && \text{Simplify} \\
 & 6 = -6 && \text{False}
 \end{aligned}$$

$x = 33$ does not check because $\sqrt{36} = 6$, not -6 . We conclude that there is *no solution* to this equation, and the solution set is \emptyset .

► **Quick check** Find the solution set. $\sqrt{x+5} = 6$ and $\sqrt{2x+5} = \sqrt{x+8}$ ■

Mastery points

Can you

- Solve equations containing radicals?

Exercise 9-7

Find the solution set. See example 9-7 A.

Examples $\sqrt{x+5} = 6$

Solutions $(\sqrt{x+5})^2 = (6)^2$
 $x+5 = 36$
 $x = 31$

Check:

$$\begin{aligned}
 \sqrt{(31)+5} &= 6 \\
 \sqrt{36} &= 6 \\
 6 &= 6
 \end{aligned}$$

True

The solution set is $\{31\}$.

$$\sqrt{2x+5} = \sqrt{x+8}$$

$$\begin{aligned}
 (\sqrt{2x+5})^2 &= (\sqrt{x+8})^2 && \text{Square both members} \\
 2x+5 &= x+8 && \text{Simplify} \\
 x+5 &= 8 && \text{Solve for } x \\
 x &= 3
 \end{aligned}$$

Check:

$$\begin{aligned}
 \sqrt{2(3)+5} &= \sqrt{(3)+8} && \text{Substitute into original equation} \\
 \sqrt{6+5} &= \sqrt{11} && \text{Simplify} \\
 \sqrt{11} &= \sqrt{11} && \text{True}
 \end{aligned}$$

The solution set is $\{3\}$.

- | | | | |
|----------------------------------|---------------------------------|----------------------------------|-------------------------|
| 1. $\sqrt{x} = 4$ | 2. $\sqrt{x} = 5$ | 3. $\sqrt{x} = 9$ | 4. $\sqrt{x} = 7$ |
| 5. $\sqrt{x+5} = 4$ | 6. $\sqrt{x-3} = 5$ | 7. $\sqrt{x-7} = 6$ | 8. $\sqrt{x+3} = 7$ |
| 9. $\sqrt{2x+1} = 5$ | 10. $\sqrt{2x+6} = 4$ | 11. $\sqrt{3x+1} = 4$ | 12. $\sqrt{5x-4} = 6$ |
| 13. $\sqrt{x} + 4 = 7$ | 14. $\sqrt{x} + 2 = 9$ | 15. $\sqrt{x} - 5 = 1$ | 16. $\sqrt{x} - 4 = 2$ |
| 17. $\sqrt{x} + 7 = 5$ | 18. $\sqrt{x} + 8 = 4$ | 19. $\sqrt{x} + 6 = 3$ | 20. $\sqrt{x} + 10 = 5$ |
| 21. $\sqrt{2x+1} = \sqrt{x+5}$ | 22. $\sqrt{2x+4} = \sqrt{3x-2}$ | 23. $\sqrt{5x-3} = \sqrt{2x+9}$ | |
| 24. $\sqrt{7x-4} = \sqrt{3x+20}$ | 25. $\sqrt{2x+7} = \sqrt{4x+1}$ | 26. $\sqrt{6x-3} = \sqrt{4x+5}$ | |
| 27. $\sqrt{3x+5} = \sqrt{5x+1}$ | 28. $\sqrt{4x-4} = \sqrt{x+5}$ | 29. $\sqrt{5x-7} = \sqrt{2x+8}$ | |
| 30. $\sqrt{4x-15} = \sqrt{7x-9}$ | 31. $\sqrt{5+2x} = \sqrt{2+3x}$ | 32. $\sqrt{9-5x} = \sqrt{15-7x}$ | |

33. $\sqrt{x}\sqrt{x-15} = 4$ 34. $\sqrt{x}\sqrt{x-3} = 2$ 35. $\sqrt{x}\sqrt{x-8} = 3$
 36. $\sqrt{x}\sqrt{x-6} = 4$ 37. $\sqrt{x}\sqrt{x+6} = 4$ 38. $\sqrt{x+3}\sqrt{x} = 2$
 39. $\sqrt{x^2+1} = x+2$ 40. $\sqrt{x^2+3x} = x+1$ 41. $\sqrt{x^2+3x} = x-3$
 42. $\sqrt{x^2+12} = x+2$ 43. $\sqrt{x+6} = x$ 44. $\sqrt{5x+6} = x$
 45. $\sqrt{2x+8} = x$ 46. $\sqrt{4x+12} = x$ 47. $\sqrt{x-4} = x-6$
 48. $\sqrt{x+2} = x+2$ 49. $\sqrt{x+4} + 8 = x$ 50. $\sqrt{x} + 6 = x$
 51. $\sqrt{x+7} = 2x-1$ 52. $\sqrt{2x-1} + 2x = 7$

Find the unknown number in problems 53–60.

53. The square root of the sum of a number and 6 is 5.
Find the number.
54. The square root of the sum of a number and 9 is 7.
Find the number.
55. The square root of the product of a number and 6 is 12. Find the number.
56. The square root of the product of a number and 9 is 6. Find the number.
57. A certain number is equal to the square root of the sum of that number and 12.
58. The square root of the product of a number and 12 is equal to the number increased by 3.
59. The square root of the sum of a number and 11 is 1 less than the number.
60. The square root of the product of a number and 4 is 3 less than the number.
61. At an altitude of h ft above the sea or level ground, the distance d in miles that a person can see an object is given by

$$d = \sqrt{\frac{3h}{2}}$$

How high must a person be to see an object 6 miles away?

62. The formula for approximating the velocity V in miles per hour of a car based on the length of its skid marks S (in feet) on dry pavement is given by
 $V = 2\sqrt{6S}$
 If the velocity is 36 mph, how long will the skid marks be?
63. On wet pavement, the formula in exercise 62 is given by
 $V = 2\sqrt{3S}$
 How long will the skid marks be if the car is traveling at 36 mph on wet pavement?
64. Find the number whose principal square root is 3.
(Hint: Let $\sqrt{x} = 3$)
65. Find the number whose principal square root is 10.
66. Find the number whose principal square root is 11.


Review exercises

Completely factor the following expressions. See sections 4–2 and 4–4.

1. $x^2 - 4$ 2. $x^2 + 9x + 18$ 3. $x^2 - 3x - 10$ 4. $x^2 - 6x + 9$

Simplify. See section 9–1.

5. $\sqrt{81}$ 6. $\sqrt{49}$ 7. $\sqrt{121}$
 8. Find the solution set for the equation
 $x^2 = 64$. See section 4–7.



Extra Credit Rocks

Sign up for a Discover® Student Card today and enjoy:

- 0% Intro APR* on Purchases for 6 Months
- No Annual Fee
- Easiest Online Account Management Options
- Full 5% *Cashback Bonus*®* on Get More purchases in popular categories all year
- Up to 1% *Cashback Bonus*®* on all your other purchases
- Unlimited cash rewards that never expire as long as you use your Card

APPLY NOW

DISCOVER
CARD

*View Discover® Card Rates, Fees, Rewards and Other Important Information.

Chapter 9 lead-in problem

Roger has a ladder that will extend to a length of 21 feet. For the ladder to be safe to climb on, it must be placed 7 feet away from the house. The roof is 20 feet above the ground. Will the ladder be able to reach the roof safely? If not, how far up will the ladder reach? (Leave your answer rounded to one decimal place.)

Solution



$$\begin{aligned}
 a^2 + b^2 &= c^2 && \text{Pythagorean Theorem} \\
 a^2 + (7)^2 &= (21)^2 && \text{Substitute 21 for } c \text{ and 7 for } b \\
 a^2 + 49 &= 441 && \text{Simplify} \\
 a^2 &= 392 && \text{Isolate } a^2 \\
 a &\approx 19.8 && \text{Round } \sqrt{392} \text{ to one decimal place}
 \end{aligned}$$

The ladder will not be able to reach the roof safely. The ladder's maximum safe reach is approximately 19.8 feet.

Chapter 9 summary

- $\sqrt[n]{a} = b$ if $\overbrace{b \cdot b \cdot b \cdots b}^{n \text{ factors}} = b^n = a$, where n is a natural number greater than 1.
- If we exclude even roots of negative numbers, which do not exist in the set of real numbers, the **principal n th root** of a number, denoted by $\sqrt[n]{}$, has the same sign as the number itself.
- When we multiply two radicals *having the same index*, we multiply the radicands and put the product under a radical symbol with the common index. $\sqrt{a}\sqrt{b} = \sqrt{ab}$ and $\sqrt[n]{a}\sqrt[n]{b} = \sqrt[n]{ab}$, **product property for radicals**.
- We can simplify a radical if the radicand has a factor(s) whose exponent is equal to or greater than the value of the index.
- The n th root of a *fraction* can be written as the n th root of the numerator over the n th root of the denominator.

$$\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}} \text{ and } \sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}, \text{ where } b \neq 0, \text{ quotient property for radicals.}$$
- We eliminate radicals from the denominator of a fraction by **rationalizing** the denominator.
- We can only add or subtract **like radicals**.
- Conjugate factors** are used to rationalize a denominator when the denominator has two terms where one or both terms contain a square root.
- When a number is raised to a **fractional exponent**, the numerator of the fractional exponent indicates the power to which the base is to be raised, and the denominator indicates the root to be taken.
- If a is any real number, m is any integer, and n is any positive integer, then if $\frac{m}{n}$ is reduced to lowest terms, $a^{m/n} = (\sqrt[n]{a})^m = \sqrt[n]{a^m}$ provided that $\sqrt[n]{a}$ is a real number.
- The **squaring property of equality** states that if both members of an equation are squared, the solution(s) of the original equation are solution(s) of the resulting equation.
- When solving equations where the unknown is under a radical symbol, we must check for **extraneous roots**.

Chapter 9 error analysis

1. Principal square root

Example: $\sqrt{36} = -6$

Correct answer: 6

What error was made? (see page 368)

2. Rationalizing the denominator of an n th root

Example: $\frac{x}{\sqrt[3]{x}} = \frac{x}{\sqrt[3]{x}} \cdot \frac{\sqrt[3]{x}}{\sqrt[3]{x}} = \frac{x\sqrt[3]{x}}{x} = \sqrt[3]{x}$

Correct answer: $\sqrt[3]{x^2}$

What error was made? (see page 380)

3. Addition in the radicand of a radical expression

Example: $\sqrt{9+4} = \sqrt{9} + \sqrt{4} = 3 + 2 = 5$

Correct answer: $\sqrt{13}$

What error was made? (see page 374)

4. Subtraction of radical expressions

Example: $\sqrt{3} - \sqrt{2} = \sqrt{3-2} = \sqrt{1} = 1$

Correct answer: $\sqrt{3} - \sqrt{2}$

What error was made? (see page 374)

5. Sums and differences of radical numbers

Example:

$\sqrt{8} + 3\sqrt{2} - \sqrt{18} = 2\sqrt{2} + 3\sqrt{2} - \sqrt{18}$
 $= 5\sqrt{2} - \sqrt{18}$

Correct answer: $2\sqrt{2}$

What error was made? (see page 384)

6. Multiplication of radical expressions

Example: $\sqrt{3}(\sqrt{3} + \sqrt{2}) = \sqrt{3} \cdot \sqrt{3} + \sqrt{2} = 3 + \sqrt{2}$

Correct answer: $3 + \sqrt{6}$

What error was made? (see page 387)

7. Squaring a radical binomial

Example: $(\sqrt{2} - \sqrt{3})^2 = (\sqrt{2})^2 - (\sqrt{3})^2 = 2 - 3 = -1$

Correct answer: $5 - 2\sqrt{6}$

What error was made? (see page 387)

8. Fractional exponents to radical expressions

Example: $x^{3/2} = \sqrt[3]{x^2}$

Correct answer: $x\sqrt{x}$

What error was made? (see page 392)

9. Extraneous solutions of radical equations

Example: Find the solution set of $\sqrt{x+2} = x-4$

$(\sqrt{x+2})^2 = (x-4)^2$

$x+2 = x^2 - 8x + 16$

$x^2 - 9x + 14 = 0$

$(x-7)(x-2) = 0$

$x-7 = 0$ or $x-2 = 0$

$x = 7$ or $x = 2$ $\{2, 7\}$

Correct answer: $\{7\}$

What error was made? (see page 396)

10. Negative exponents

Example: $6^{-2} = -36$

Correct answer: $\frac{1}{36}$

What error was made? (see page 141)

Chapter 9 critical thinking

Given the numbers 33 and 27, determine a method by which you can multiply these numbers mentally.

Chapter 9 review

Assume that all variables in problems 1–42 represent positive real numbers and that no denominator is equal to zero.

[9–1]

Find the indicated root.

1. $\sqrt{81}$

2. $\sqrt{25}$

3. $-\sqrt{9}$

4. $-\sqrt{49}$

[9–2]

Perform any indicated operations and simplify.

5. $\sqrt{40}$

6. $\sqrt{18a^2b^3}$

7. $\sqrt{2}\sqrt{14}$

8. $\sqrt{18}\sqrt{10}$

[9–3]

Express the given radicals in simplest form with all denominators rationalized.

9. $\sqrt{\frac{16}{17}}$

10. $\sqrt{\frac{7}{18}}$

11. $\sqrt{\frac{a}{b}}$

12. $\sqrt{\frac{x}{y^3}}$

13. $\frac{a}{\sqrt{ab}}$

14. $\frac{2x}{\sqrt{xy}}$

[9-4]

Perform the indicated operations and simplify.

15. $3\sqrt{7} + 4\sqrt{7}$

16. $\sqrt{18} + 5\sqrt{2}$

17. $3\sqrt{20} - \sqrt{45}$

18. $2\sqrt{75} - \sqrt{3} + 5\sqrt{27}$

19. $\sqrt{50a} - 2\sqrt{8a}$

20. $7\sqrt{9x} - 2\sqrt{4x}$

[9-5]

Perform the indicated operations and simplify.

21. $\sqrt{3}(\sqrt{5} - \sqrt{7})$

22. $\sqrt{10}(\sqrt{14} + \sqrt{6})$

23. $(5 + \sqrt{7})(3 - \sqrt{7})$

24. $(6 - \sqrt{3})^2$

25. $(\sqrt{3} + \sqrt{5})^2$

26. $(2\sqrt{a} - \sqrt{b})(2\sqrt{a} + \sqrt{b})$

Express the given radicals in simplest form with all denominators rationalized.

27. $\frac{1}{\sqrt{3} - 2}$

28. $\frac{2}{\sqrt{6} + 4}$

29. $\frac{1}{\sqrt{a} + b}$

30. $\frac{x}{\sqrt{xy} + x}$

31. $\frac{a}{a + \sqrt{b}}$

32. $\frac{\sqrt{2} + 3}{\sqrt{2} - 3}$

[9-6]

Simplify the given expressions.

33. $(36)^{1/2}$

34. $(8)^{2/3}$

35. $(-8)^{1/3}$

36. $(32)^{-2/5}$

Perform the indicated operations and simplify.

37. $a^{2/5} \cdot a^{3/5}$

38. $b^{1/3} \cdot b^{3/4}$

39. $\frac{a^{1/2}}{a^{1/4}}$

40. $(a^{3/2})^{3/4}$

41. $(16a^4b^8)^{3/4}$

42. $\frac{a^2b^2}{a^{1/2}b^{3/2}}$

[9-7]

Find the solution set.

43. $\sqrt{x} = 8$

44. $\sqrt{x - 4} = 7$

45. $\sqrt{5x - 3} = \sqrt{3x + 5}$

46. $\sqrt{x}\sqrt{x + 8} = 3$

47. $\sqrt{x}\sqrt{x + 6} = 4$

48. $\sqrt{x^2 + 16} = x + 2$

49. $\sqrt{x + 6} = x + 4$

50. $\sqrt{x - 3} = x - 3$

Chapter 9 cumulative test

Perform the indicated operations and simplify. Assume that all variables represent positive real numbers and that no denominator is equal to zero.

[1-8] 1. $3[4(6 - 2) + (-5 + 4)]$

[3-1] 2. $x^3 \cdot x^2 \cdot x^2$

[3-3] 3. $x^{12} \div x^6$

[2-3] 4. $(3x^2 - 2x + 5) - (x^2 - 4x - 8)$

[3-2] 5. $4a^2b^3(2a^2 - 3ab + 4b^2)$

[3-3] 6. $\frac{8a^{-2}b^3c^0}{4a^{-5}b}$

[1-8] 7. -6^2

[3-2] 8. $(3a - b)^2$

[6-1] 9. $\frac{a^2 - 9}{3a + 12} \cdot \frac{a + 4}{a^2 + 5a + 6}$

[9-4] 10. $5\sqrt{12} + 2\sqrt{27}$

[9-1] 11. $\sqrt[3]{-8}$

[6-2] 12. $\frac{2x}{x - 1} - \frac{x - 3}{x - 1}$

[9-5] 13. $\frac{\sqrt{x}}{x - \sqrt{y}}$

[9-2] 14. $\sqrt[3]{81x^4y^6z}$

[3-2] 15. $(5x - y)(5x + y)$

[2-3] 16. $3x - [2x - (x - y) + 3y]$

Factor completely.

[4-1] 17. $6a^3b^4 - 2a^3b^5 + 8a^5b^3$

[4-3] 19. $2x^2 + 7x - 4$

[4-3] 21. $6x^2 + 11x + 4$

[4-4] 18. $25c^2 - d^2$

[4-4] 20. $y^4 - 4z^2$

[4-2] 22. $x^2 + 3x - 28$

Find the solution set for problems 23–28 and solve problems 29 and 30.

[2-6] 23. $2(4x - 3) = 5x - 7$

[2-6] 25. $3(x + 4) - 2(x - 3) = 12$

[6-5] 27. $\frac{3x + 1}{9} + \frac{1}{12} = \frac{2x - 1}{3}$

[2-9] 29. $-3 < 2x - 5 < 11$

[7-3] 31. Find the slope of the line passing through points (4,2) and (3,5).

[8-2] 33. Solve the system of equations.
 $5x - y = 4$
 $x + 3y = 2$

[5-4] 35. A punch machine can make 18 holes in 4 minutes. How many holes can the machine make in 5 hours?

[2-8] 37. The width of a rectangle is 6 feet less than its length. The perimeter of the rectangle is 96 feet. Find the dimensions.

[4-7] 24. $x^2 = 9$

[2-6] 26. $\frac{1}{4}x + 2 = \frac{1}{2}x - 1$

[4-7] 28. $2x^2 + 3x + 1 = 0$

[2-9] 30. $3x + 5 > x + 12$

[7-4] 32. Write $8x - 2y = 4$ in slope-intercept form and determine the slope and y-intercept.

[2-8] 34. One number is four times a second number and their sum is 70. Find the numbers.

[4-8] 36. The product of two consecutive positive even integers is 288. Find the integers.

Chapter 9

Exercise 9-1

Answers to odd-numbered problems

1. 10 3. 2 5. -12 7. -11 9. 11 11. -4 13. 4.243
15. 6.403 17. -7.211 19. 24 amperes 21. 13 units
23. 5 meters 25. 13 inches 27. 8 yards 29. 20 millimeters
31. 2 33. 5 35. -2 37. 3 39. -3 41. 2 43. -3
45. 1 47. -1 49. 9 units

Solutions to trial exercise problems

$$19. I = \sqrt{\frac{\text{watts}}{\text{ohms}}} = \sqrt{\frac{(\quad)}{(\quad)}} = \sqrt{\frac{1,728}{3}} = \sqrt{576} = 24$$

Answer: 24 amperes 31. $\sqrt[3]{8} = 2$, since $2 \cdot 2 \cdot 2 = 2^3 = 8$.
39. $-\sqrt[4]{81} = -3$, since $3 \cdot 3 \cdot 3 \cdot 3 = 3^4 = 81$ and the negative sign indicates that we want the negative root. 45. $\sqrt[10]{1} = 1$, since $1^{10} = 1$. 1 raised to any power is 1 and the n th root of 1 is 1.

Review exercises

1. 3^2 2. $2^2 \cdot 3$ 3. 2^3 4. $2^3 \cdot 5$ 5. $2 \cdot 5^2$ 6. 3^4 7. 2^6
8. 2^4

Exercise 9-2

Answers to odd-numbered problems

1. 4 3. $2\sqrt{5}$ 5. $3\sqrt{5}$ 7. $4\sqrt{2}$ 9. $4\sqrt{3}$ 11. $7\sqrt{2}$
13. $a^3\sqrt{a}$ 15. $2ab\sqrt{b}$ 17. $3ab^2\sqrt{3ab}$ 19. $3\sqrt{2}$ 21. 15
23. $2\sqrt{15}$ 25. $5\sqrt{15}$ 27. $5\sqrt{3}$ 29. $5x\sqrt{3}$ 31. $4b\sqrt{3a}$
33. 13 feet 35. 24 mph 37. $2\sqrt[5]{2}$ 39. $2\sqrt[3]{3}$ 41. $b^2\sqrt[3]{b^2}$
43. y^3 45. $b\sqrt[3]{4a^2}$ 47. $2ab\sqrt[3]{2ab^2}$ 49. $3ab^3\sqrt[3]{3a^2b^2}$
51. $b\sqrt[3]{b}$ 53. a 55. $3ab\sqrt[3]{2b}$ 57. $3b\sqrt[3]{3a^3}$ 59. $4a^5b^3\sqrt[3]{3b}$
61. 2 inches 63. 2 inches

Solutions to trial exercise problems

$$15. \sqrt{4a^2b^3} = \sqrt{4 \cdot a^2 \cdot b^2 \cdot b} = \sqrt{4} \sqrt{a^2} \sqrt{b^2} \sqrt{b} = 2ab\sqrt{b}$$

$$19. \sqrt[3]{6\sqrt{3}} = \sqrt[3]{18} = \sqrt[3]{9 \cdot 2} = \sqrt[3]{9} \sqrt[3]{2} = 3\sqrt[3]{2}$$

$$42. \sqrt[3]{x^9} = \sqrt[3]{x^3 \cdot x^3 \cdot x^3} = \sqrt[3]{x^3} \sqrt[3]{x^3} \sqrt[3]{x^3} = x \cdot x \cdot x = x^3$$

$$54. \sqrt[3]{5a^2b} \sqrt[3]{75a^2b^2} = \sqrt[3]{375a^4b^3} = \sqrt[3]{125 \cdot 3 \cdot a^3 \cdot a \cdot b^3}$$

$$= \sqrt[3]{125} \sqrt[3]{a^3} \sqrt[3]{b^3} \sqrt[3]{3a} = 5ab\sqrt[3]{3a}$$

$$61. h = \sqrt[3]{\frac{12I}{b}} = \sqrt[3]{\frac{12(\quad)}{(\quad)}} = \sqrt[3]{\frac{12(2)}{3}} = \sqrt[3]{\frac{24}{3}} = \sqrt[3]{8} = 2$$

Answer: $h = 2$ inches

Review exercises

1. $\frac{7}{8}$ 2. $-4x^2y$ 3. $\frac{2y+10}{y+1}$ 4. 5 5. 3 6. 4 7. 6
8. x

Exercise 9-3

Answers to odd-numbered problems

1. $\frac{3}{5}$ 3. $\frac{5}{7}$ 5. $\frac{\sqrt{3}}{2}$ 7. $\frac{8}{a}$ 9. $\frac{\sqrt{2}}{2}$ 11. $\frac{2\sqrt{7}}{7}$ 13. $\frac{\sqrt{15}}{15}$
15. $\frac{2\sqrt{3}}{15}$ 17. $\sqrt{2}$ 19. $\frac{5\sqrt{2}}{2}$ 21. $\frac{x\sqrt{y}}{y}$ 23. $\frac{\sqrt{x}}{x}$ 25. a^2
27. $\sqrt{41}$ meters 29. 12 feet 31. $\frac{1}{2}$ 33. $\frac{3}{5}$ 35. $\frac{a^2\sqrt[3]{3}}{b}$
37. $\frac{\sqrt[5]{a^4}}{b^2}$ 39. $\frac{ab^2\sqrt[4]{bc}}{c^3}$ 41. $\frac{\sqrt[5]{x^3y^2}}{z^3}$ 43. $\frac{\sqrt[3]{20}}{5}$ 45. $\frac{2\sqrt[4]{5}}{5}$
47. $\frac{x\sqrt[3]{y}}{y}$ 49. $b\sqrt[3]{a}$ 51. $\frac{a\sqrt[3]{bc^2}}{bc}$ 53. $\frac{\sqrt[3]{a^2bc^2}}{bc}$ 55. $\sqrt[3]{a^2b}$
57. 3 units

Solutions to trial exercise problems

$$15. \sqrt{\frac{4}{75}} = \frac{\sqrt{4}}{\sqrt{75}} = \frac{\sqrt{2^2}}{\sqrt{3 \cdot 5^2}} = \frac{2}{5\sqrt{3}} \cdot \frac{\sqrt{3}}{\sqrt{3}} = \frac{2\sqrt{3}}{5 \cdot 3} = \frac{2\sqrt{3}}{15}$$

$$35. \sqrt[3]{\frac{3a^6}{b^3}} = \frac{\sqrt[3]{3a^6}}{\sqrt[3]{b^3}} = \frac{a^2\sqrt[3]{3}}{b}$$

$$39. \sqrt[4]{\frac{a^4b^9}{c^{11}}} = \frac{\sqrt[4]{a^4b^9}}{\sqrt[4]{c^{11}}} = \frac{ab^2\sqrt[4]{b}}{c^2\sqrt[4]{c^3}} = \frac{ab^2\sqrt[4]{b}}{c^2\sqrt[4]{c^3}}$$

$$45. \sqrt[4]{\frac{16}{125}} = \frac{\sqrt[4]{16}}{\sqrt[4]{125}} = \frac{\sqrt[4]{2^4}}{\sqrt[4]{5^3}} = \frac{2}{\sqrt[4]{5^3}} \cdot \frac{\sqrt[4]{5}}{\sqrt[4]{5}} = \frac{2\sqrt[4]{5}}{\sqrt[4]{5^4}} = \frac{2\sqrt[4]{5}}{5}$$

$$49. \frac{ab}{\sqrt[3]{a^2}} \cdot \frac{\sqrt[3]{a^1}}{\sqrt[3]{a^1}} = \frac{ab\sqrt[3]{a}}{\sqrt[3]{a^3}} = \frac{ab\sqrt[3]{a}}{a} = b\sqrt[3]{a}$$

$$51. \sqrt[3]{\frac{a^3}{b^2c}} = \frac{\sqrt[3]{a^3}}{\sqrt[3]{b^2c}} = \frac{a}{\sqrt[3]{b^2c}} \cdot \frac{\sqrt[3]{b^1c^2}}{\sqrt[3]{b^1c^2}} = \frac{a\sqrt[3]{bc^2}}{\sqrt[3]{b^3c^3}} = \frac{a\sqrt[3]{bc^2}}{bc}$$

$$57. r = \sqrt[3]{\frac{3V}{4\pi}} = \sqrt[3]{\frac{3(\quad)}{4(\quad)}} = \sqrt[3]{\frac{3(113.04)}{4(3.14)}} = \sqrt[3]{\frac{339.12}{12.56}}$$

$$= \sqrt[3]{27} = \sqrt[3]{3^3} = 3 \quad \text{Answer: 3 units}$$

Review exercises

1. $6x$ 2. $4y$ 3. $8ab$ 4. $5xy$ 5. $x^2 - 9$ 6. $x^2 - y^2$

Exercise 9-4

Answers to odd-numbered problems

1. $9\sqrt{3}$ 3. $10\sqrt{5}$ 5. $6\sqrt{3}$ 7. $3\sqrt{7}$ 9. $3\sqrt{a}$ 11. $8\sqrt{a}$
13. $7\sqrt{xy}$ 15. $8\sqrt{a} + 2\sqrt{ab}$ 17. $4\sqrt{xy} + 3\sqrt{y}$ 19. $7\sqrt{2}$
21. $\sqrt{3}$ 23. $17\sqrt{7}$ 25. $7\sqrt{2}$ 27. $2\sqrt{3} + 8\sqrt{2}$ 29. $7\sqrt{2a}$
31. $-\sqrt{x}$ 33. $17\sqrt{2a}$ 35. $2\sqrt{2a} + 6\sqrt{3a}$ 37. 14 units
39. 8.1 units by 8.1 units 41. $6\sqrt[5]{2}$ 43. $5\sqrt[3]{2}$
45. $3\sqrt[3]{3} + 10\sqrt[3]{2}$ 47. $5\sqrt[4]{x^3}$ 49. $\sqrt[3]{x^2y}$ 51. $2a\sqrt[3]{b^2}$

Solutions to trial exercise problems

$$14. 3\sqrt{x} + 2\sqrt{y} - \sqrt{x} = 3\sqrt{x} - \sqrt{x} + 2\sqrt{y} = 2\sqrt{x} + 2\sqrt{y}$$

$$18. \sqrt{20} + 3\sqrt{5} = \sqrt{2^2 \cdot 5} + 3\sqrt{5} = 2\sqrt{5} + 3\sqrt{5} = 5\sqrt{5}$$

$$29. \sqrt{50a} + \sqrt{8a} = \sqrt{2 \cdot 5^2 \cdot a} + \sqrt{2^3 \cdot a} = 5\sqrt{2a} + 2\sqrt{2a}$$

$$= 7\sqrt{2a} \quad 37. h = b + s, \text{ where } s = 6 \text{ units and } b = \sqrt{c^2 - s^2}$$

$$= \sqrt{(10)^2 - (6)^2} = \sqrt{100 - 36} = \sqrt{64} = 8. \text{ Then } h = b + s$$

$$= (8) + (6) = 14. \text{ Answer: 14 units}$$

$$43. \sqrt[3]{16} + \sqrt[3]{54} = \sqrt[3]{2^4} + \sqrt[3]{2 \cdot 3^3} = 2\sqrt[3]{2} + 3\sqrt[3]{2} = 5\sqrt[3]{2}$$

$$50. \sqrt[3]{x^6y} + 2x^2\sqrt[3]{y} = x^2\sqrt[3]{y} + 2x^2\sqrt[3]{y} = (1 + 2)x^2\sqrt[3]{y} = 3x^2\sqrt[3]{y}$$

Review exercises

1. $6x^2 - 3xy$ 2. $2a^4 - 2a^2b^2$ 3. $x^2 - 2x + 1$ 4. $y^2 - 1$
5. $4x^2 - 1$ 6. $x^2 + 5xy + 6y^2$ 7. $x^2 - 2xy + y^2$
8. $a^2 + 4ab + 4b^2$

Exercise 9-5

Answers to odd-numbered problems

1. $3\sqrt{2} + 3\sqrt{3}$ 3. $\sqrt{6} + \sqrt{14}$ 5. $6\sqrt{6} - 3\sqrt{22}$
7. $5\sqrt{3} - 5\sqrt{2}$ 9. $14\sqrt{5} - 42\sqrt{2}$ 11. $3a + \sqrt{ab}$
13. $14 + 7\sqrt{2}$ 15. $12 - 25\sqrt{a} + 12a$ 17. 1 19. -2
21. $9 + 4\sqrt{5}$ 23. $x + 2\sqrt{xy} + y$ 25. $x - y$ 27. $x^2y - z$
29. $4x + 4y\sqrt{x} + y^2$ 31. $11 + \sqrt{3}$ 33. $\sqrt{a} - 3\sqrt{b}$
35. $\frac{-\sqrt{2} + 3}{7}$ 37. $\frac{-14 + 7\sqrt{7}}{3}$ 39. $\sqrt{6} + \sqrt{3}$
41. $\frac{6\sqrt{3} + 3\sqrt{5}}{7}$ 43. $\frac{-3 - \sqrt{5}}{2}$ 45. $\frac{a + 2b\sqrt{a} + b^2}{a - b^2}$

Solutions to trial exercise problems

$$\begin{aligned}
 5. & 3\sqrt{2}(2\sqrt{3} - \sqrt{11}) = 3\sqrt{2} \cdot 2\sqrt{3} - 3\sqrt{2} \cdot \sqrt{11} \\
 & = 6\sqrt{6} - 3\sqrt{22} \quad 15. (3 - 4\sqrt{a})(4 - 3\sqrt{a}) \\
 & = 3 \cdot 4 - 3 \cdot 3\sqrt{a} - 4\sqrt{a} \cdot 4 + 4\sqrt{a} \cdot 3\sqrt{a} \\
 & = 12 - 9\sqrt{a} - 16\sqrt{a} + 12a = 12 - 25\sqrt{a} + 12a \\
 17. & (\sqrt{3} + \sqrt{2})(\sqrt{3} - \sqrt{2}) \text{ Conjugates, therefore} \\
 & = (\sqrt{3})^2 - (\sqrt{2})^2 = 3 - 2 = 1 \quad 21. (2 + \sqrt{5})^2 \\
 & = (2 + \sqrt{5})(2 + \sqrt{5}) = 2 \cdot 2 + 2\sqrt{5} + 2\sqrt{5} + \sqrt{5}\sqrt{5} \\
 & = 4 + 4\sqrt{5} + 5 = 9 + 4\sqrt{5} \quad 26. (2\sqrt{a} - \sqrt{b})(2\sqrt{a} + \sqrt{b}) \\
 & \text{Conjugates, therefore} = (2\sqrt{a})^2 - (\sqrt{b})^2 = 2^2(\sqrt{a})^2 - b \\
 & = 4a - b \quad 38. \frac{6}{3 - \sqrt{6}} = \frac{6}{3 - \sqrt{6}} \cdot \frac{3 + \sqrt{6}}{3 + \sqrt{6}} = \frac{6(3 + \sqrt{6})}{(3)^2 - (\sqrt{6})^2} \\
 & = \frac{6(3 + \sqrt{6})}{9 - 6} = \frac{6(3 + \sqrt{6})}{3} = 2(3 + \sqrt{6}) = 6 + 2\sqrt{6} \\
 43. & \frac{1 + \sqrt{5}}{1 - \sqrt{5}} = \frac{1 + \sqrt{5}}{1 - \sqrt{5}} \cdot \frac{1 + \sqrt{5}}{1 + \sqrt{5}} = \frac{(1)^2 + \sqrt{5} + \sqrt{5} + (\sqrt{5})^2}{(1)^2 - (\sqrt{5})^2} \\
 & = \frac{1 + 2\sqrt{5} + 5}{1 - 5} = \frac{6 + 2\sqrt{5}}{-4} = \frac{2(3 + \sqrt{5})}{-4} \\
 & = \frac{1 - 5}{-4} = \frac{-4}{-4} = 1 \\
 & = \frac{-(3 + \sqrt{5})}{2} = \frac{-3 - \sqrt{5}}{2}
 \end{aligned}$$

Review exercises

$$\begin{aligned}
 1. & 64 \quad 2. 32 \quad 3. 9 \quad 4. \frac{1}{27} \quad 5. \frac{1}{x^6} \quad 6. 8a^6b^3 \\
 7. & \frac{y^3}{x} \quad 8. \frac{1}{x^9}
 \end{aligned}$$

Exercise 9-6

Answers to odd-numbered problems

$$\begin{aligned}
 1. & 6 \quad 3. a^2 \quad 5. 2 \quad 7. -3 \quad 9. 9 \quad 11. 27 \quad 13. \frac{1}{5} \quad 15. \frac{1}{8} \\
 17. & -\frac{1}{2} \quad 19. 4 \quad 21. 2^{5/6} \quad 23. x \quad 25. c^{3/4} \quad 27. 2 \quad 29. 2^{1/6} \\
 31. & a^{3/5} \quad 33. y^{1/6} \quad 35. a^{1/3} \quad 37. x^{2/3} \quad 39. c^{1/6} \quad 41. a^{1/3} \\
 43. & \frac{1}{x^{1/6}} = \frac{x^{5/6}}{x} \quad 45. \frac{1}{y^{1/4}} = \frac{y^{3/4}}{y} \quad 47. 8a^3 \quad 49. 4a^4b^2 \\
 51. & b^{1/2}c^{1/4} \quad 53. a^{1/2}b^{2/3} \quad 55. 81 \quad 57. 49 \text{ mph} \quad 59. 24 \text{ miles} \\
 61. & \text{Take the square root 3 times, that is, the square root of the square root of the square root is the eighth root.}
 \end{aligned}$$

Solutions to trial exercise problems

$$\begin{aligned}
 3. & (a^6)^{1/3} = a^{6/1 \cdot 1/3} = a^2 \quad 7. (-27)^{1/3} = \sqrt[3]{(-27)} = -3 \\
 9. & (27)^{2/3} = (\sqrt[3]{27})^2 = (3)^2 = 9 \quad 17. (-8)^{-1/3} = \frac{1}{(-8)^{1/3}} \\
 & = \frac{1}{\sqrt[3]{-8}} = \frac{1}{-2} = -\frac{1}{2} \quad 35. (a^{2/3})^{1/2} = a^{2/3 \cdot 1/2} = a^{1/3} \\
 39. & (c^{-1/4})^{-2/3} = c^{(-1/4) \cdot (-2/3)} = c^{1/6} \quad 51. \frac{b^{3/4}c^{1/2}}{b^{1/4}c^{1/4}} \\
 & = b^{3/4 - 1/4}c^{1/2 - 1/4} = b^{2/4}c^{2/4 - 1/4} = b^{1/2}c^{1/4}
 \end{aligned}$$

Review exercises

$$\begin{aligned}
 1. & 7 \quad 2. x \quad 3. x + 1 \quad 4. x^2 + 2x + 1 \quad 5. x^2 - 4x + 4 \\
 6. & \{-2, 3\} \quad 7. \{2, 8\} \quad 8. \{-1, 0\}
 \end{aligned}$$

Exercise 9-7

Answers to odd-numbered problems

$$\begin{aligned}
 1. & \{16\} \quad 3. \{81\} \quad 5. \{11\} \quad 7. \{43\} \quad 9. \{12\} \quad 11. \{5\} \\
 13. & \{9\} \quad 15. \{36\} \quad 17. \emptyset \quad 19. \emptyset \quad 21. \{4\} \quad 23. \{4\} \quad 25. \{3\} \\
 27. & \{2\} \quad 29. \{5\} \quad 31. \{3\} \quad 33. \{16\} \quad 35. \{9\} \quad 37. \{2\}
 \end{aligned}$$

$$\begin{aligned}
 39. & \left\{-\frac{3}{4}\right\} \quad 41. \emptyset \quad 43. \{3\} \quad 45. \{4\} \quad 47. \{8\} \quad 49. \{12\} \\
 51. & \{2\} \quad 53. 19 \quad 55. 24 \quad 57. 4 \quad 59. 5 \quad 61. 24 \text{ feet} \\
 63. & 108 \text{ feet} \quad 65. 100
 \end{aligned}$$

Solutions to trial exercise problems

$$\begin{aligned}
 17. & \sqrt{x} + 7 = 5 \\
 & \sqrt{x} = -2 \\
 & (\sqrt{x})^2 = (-2)^2 \\
 & x = 4
 \end{aligned}$$

Check:

$$\begin{aligned}
 \sqrt{4} + 7 &= 5 \\
 2 + 7 &= 5 \\
 9 &= 5 \text{ (false)}
 \end{aligned}$$

Therefore no solution

and the solution set is \emptyset .

$$\begin{aligned}
 21. & \sqrt{2x+1} = \sqrt{x+5} \\
 & (\sqrt{2x+1})^2 = (\sqrt{x+5})^2 \\
 & 2x+1 = x+5 \\
 & x+1 = 5 \\
 & x = 4
 \end{aligned}$$

Check:

$$\begin{aligned}
 \sqrt{2(4)+1} &= \sqrt{(4)+5} \\
 \sqrt{8+1} &= \sqrt{9} \\
 \sqrt{9} &= \sqrt{9} \\
 3 &= 3 \text{ (true)}
 \end{aligned}$$

$$\begin{aligned}
 39. & \{4\} \\
 & \sqrt{x^2+1} = x+2 \\
 & (\sqrt{x^2+1})^2 = (x+2)^2 \\
 & x^2+1 = (x+2)(x+2) \\
 & x^2+1 = x^2+2x+2x+4 \\
 & x^2+1 = x^2+4x+4 \\
 & 1 = 4x+4 \\
 & -3 = 4x \\
 & -\frac{3}{4} = x
 \end{aligned}$$

Check:

$$\begin{aligned}
 \sqrt{\left(-\frac{3}{4}\right)^2+1} &= \left(-\frac{3}{4}\right)+2 \\
 \sqrt{\frac{9}{16}+1} &= -\frac{3}{4}+\frac{8}{4} \\
 \sqrt{\frac{9}{16}+\frac{16}{16}} &= \frac{5}{4} \\
 \sqrt{\frac{25}{16}} &= \frac{5}{4} \\
 \frac{\sqrt{25}}{\sqrt{16}} &= \frac{5}{4} \\
 \frac{5}{4} &= \frac{5}{4} \text{ (true)}
 \end{aligned}$$

$$\begin{aligned}
 43. & \left\{-\frac{3}{4}\right\} \\
 & \sqrt{x+6} = x \\
 & (\sqrt{x+6})^2 = (x)^2 \\
 & x+6 = x^2 \\
 & 0 = x^2 - x - 6 \\
 & 0 = (x-3)(x+2) \\
 & x-3 = 0 \text{ or } x+2 = 0 \\
 & x = 3 \text{ or } x = -2
 \end{aligned}$$

Check:

$$\begin{aligned}\sqrt{(3)} + 6 &= (3) \\ \sqrt{9} &= 3 \\ 3 &= 3 \text{ (true)} \\ \sqrt{(-2)} + 6 &= (-2) \\ \sqrt{4} &= -2 \\ 2 &= -2 \text{ (false)}\end{aligned}$$

 $\{3\}$

$$\begin{aligned}47. \quad \sqrt{x-4} &= x-6 \\ (\sqrt{x-4})^2 &= (x-6)^2 \\ x-4 &= (x-6)(x-6) \\ x-4 &= x^2-6x-6x+36 \\ x-4 &= x^2-12x+36 \\ 0 &= x^2-13x+40 \\ 0 &= (x-8)(x-5) \\ x-8 &= 0 \text{ or } x-5 = 0 \\ x &= 8 \text{ or } x = 5\end{aligned}$$

Check:

$$\begin{aligned}\sqrt{(8)} - 4 &= (8) - 6 \\ \sqrt{4} &= 2 \\ 2 &= 2 \text{ (true)} \\ \sqrt{(5)} - 4 &= (5) - 6 \\ \sqrt{1} &= -1 \\ 1 &= -1 \text{ (false)}\end{aligned}$$

 $\{8\}$

 57. Let x = the number.

$$\begin{aligned}\sqrt{x+12} &= x \\ (\sqrt{x+12})^2 &= (x)^2 \\ x+12 &= x^2 \\ 0 &= x^2-x-12 \\ 0 &= (x-4)(x+3) \\ x-4 &= 0 \text{ or } x+3 = 0 \\ x &= 4 \text{ or } x = -3\end{aligned}$$

Check:

$$\begin{aligned}\sqrt{(4)} + 12 &= (4) \\ \sqrt{16} &= 4 \\ 4 &= 4 \text{ (true)} \\ \sqrt{(-3)} + 12 &= (-3) \\ \sqrt{9} &= -3 \\ 3 &= -3 \text{ (false)}\end{aligned}$$

Hence the number is 4.

Review exercises

$$\begin{aligned}1. (x+2)(x-2) \quad 2. (x+3)(x+6) \quad 3. (x+2)(x-5) \\ 4. (x-3)^2 \quad 5. 9 \quad 6. 7 \quad 7. 11 \quad 8. \{-8, 8\}\end{aligned}$$

Chapter 9 review

$$\begin{aligned}1. 9 \quad 2. 5 \quad 3. -3 \quad 4. -7 \quad 5. 2\sqrt{10} \quad 6. 3ab\sqrt{2b} \\ 7. 2\sqrt{7} \quad 8. 6\sqrt{5} \quad 9. \frac{4\sqrt{17}}{17} \quad 10. \frac{\sqrt{14}}{6} \quad 11. \frac{\sqrt{ab}}{b} \quad 12. \frac{\sqrt{xy}}{y^2} \\ 13. \frac{\sqrt{ab}}{b} \quad 14. \frac{2\sqrt{xy}}{y} \quad 15. 7\sqrt{7} \quad 16. 8\sqrt{2} \quad 17. 3\sqrt{5} \\ 18. 24\sqrt{3} \quad 19. \sqrt{2a} \quad 20. 17\sqrt{x} \quad 21. \sqrt{15} - \sqrt{21} \\ 22. 2\sqrt{35} + 2\sqrt{15} \quad 23. 8 - 2\sqrt{7} \quad 24. 39 - 12\sqrt{3} \\ 25. 8 + 2\sqrt{15} \quad 26. 4a - b \quad 27. -\sqrt{3} - 2 \quad 28. \frac{-\sqrt{6} + 4}{5} \\ 29. \frac{\sqrt{a} - b}{a - b^2} \quad 30. \frac{\sqrt{xy} - x}{y - x} \quad 31. \frac{a^2 - a\sqrt{b}}{a^2 - b} \quad 32. \frac{-11 - 6\sqrt{2}}{7}\end{aligned}$$

$$\begin{aligned}33. 6 \quad 34. 4 \quad 35. -2 \quad 36. \frac{1}{4} \quad 37. a \quad 38. b^{13/12} \quad 39. a^{1/4} \\ 40. a^{9/8} \quad 41. 8a^3b^6 \quad 42. a^{3/2}b^{1/2} \quad 43. \{64\} \quad 44. \{53\} \\ 45. \{4\} \quad 46. \{1\} \quad 47. \{2\} \quad 48. \{3\} \quad 49. \{-2\} \quad 50. \{4, 3\}\end{aligned}$$

Chapter 9 cumulative test

$$\begin{aligned}1. 45 \quad 2. x^7 \quad 3. x^6 \quad 4. 2x^2 + 2x + 13 \quad 5. 8a^4b^3 - 12a^3b^4 \\ + 16a^2b^5 \quad 6. 2a^3b^2 \quad 7. -36 \quad 8. 9a^2 - 6ab + b^2 \\ 9. \frac{a-3}{3a+6} \quad 10. 16\sqrt{3} \quad 11. -2 \quad 12. \frac{x+3}{x-1} \quad 13. \frac{x\sqrt{x} + \sqrt{xy}}{x^2 - y} \\ 14. 3xy^2\sqrt[3]{3xz} \quad 15. 25x^2 - y^2 \quad 16. 2x - 4y \\ 17. 2a^3b^3(3b - b^2 + 4a^2) \quad 18. (5c + d)(5c - d) \\ 19. (2x - 1)(x + 4) \quad 20. (y^2 + 2z)(y^2 - 2z) \\ 21. (2x + 1)(3x + 4) \quad 22. (x + 7)(x - 4) \quad 23. \left\{-\frac{1}{3}\right\} \\ 24. \{-3, 3\} \quad 25. \{-6\} \quad 26. \{12\} \quad 27. \left\{\frac{19}{12}\right\} \\ 28. \left\{-1, -\frac{1}{2}\right\} \quad 29. 1 < x < 8 \quad 30. x > \frac{7}{2} \quad 31. -3 \\ 32. y = 4x - 2; \text{ slope is 4; } y\text{-intercept is } (0, -2) \quad 33. \left(\frac{7}{8}, \frac{3}{8}\right) \\ 34. 14, 56 \quad 35. 1,350 \quad 36. 16, 18 \quad 37. 21 \text{ feet by } 27 \text{ feet}\end{aligned}$$

Chapter 10

Exercise 10–1

Answers to odd-numbered problems

$$\begin{aligned}1. \{-5, 3\} \quad 3. \left\{-\frac{3}{2}, 2\right\} \quad 5. \{-2, 2\} \quad 7. \{-8, 8\} \\ 9. \{-\sqrt{11}, \sqrt{11}\} \quad 11. \{2\sqrt{5}, -2\sqrt{5}\} \quad 13. \{-\sqrt{3}, \sqrt{3}\} \\ 15. \{-4\sqrt{2}, 4\sqrt{2}\} \quad 17. \{-3, 3\} \quad 19. \{-\sqrt{6}, \sqrt{6}\} \\ 21. \{-5\sqrt{2}, 5\sqrt{2}\} \quad 23. \{-2\sqrt{2}, 2\sqrt{2}\} \quad 25. \{-2\sqrt{2}, 2\sqrt{2}\} \\ 27. \{-\sqrt{2}, \sqrt{2}\} \quad 29. \left\{-\frac{\sqrt{6}}{5}, \frac{\sqrt{6}}{5}\right\} \quad 31. \{-\sqrt{11}, \sqrt{11}\} \\ 33. \{-4, 0\} \quad 35. \{-1, 9\} \quad 37. \{-3 - \sqrt{6}, -3 + \sqrt{6}\} \\ 39. \{9 - 3\sqrt{2}, 9 + 3\sqrt{2}\} \quad 41. \{-5 + 4\sqrt{2}, -5 - 4\sqrt{2}\} \\ 43. \{-a - 6, -a + 6\} \quad 45. \{6 - a, 6 + a\} \quad 47. \{p - q, p + q\} \\ 49. \left\{-\frac{1}{2}, \frac{7}{2}\right\} \quad 51. 5 \text{ meters} \quad 53. 2 \text{ feet} \quad 55. 9, -9 \quad 57. 0, 9 \\ 59. 7 \text{ inches, } 14 \text{ inches} \quad 61. \text{ length} = 24 \text{ meters;} \\ \text{width} = 6 \text{ meters} \quad 63. 4 \text{ and } 8 \quad 65. 5\sqrt{2} \text{ centimeters}\end{aligned}$$

Solutions to trial exercise problems

$$11. a^2 = 20$$

Extract the roots.

$$a = \sqrt{20} \text{ or } a = -\sqrt{20}$$

$$a = \sqrt{4 \cdot 5} \text{ or } a = -\sqrt{4 \cdot 5}$$

$$a = 2\sqrt{5} \text{ or } a = -2\sqrt{5}$$

$$\{2\sqrt{5}, -2\sqrt{5}\}$$

$$18. 5x^2 = 75$$

Divide each member by 5.

$$x^2 = 15$$

Extract the roots.

$$x = \sqrt{15} \text{ or } x = -\sqrt{15}$$

$$\{\sqrt{15}, -\sqrt{15}\}$$

Student Loans for up to **\$40,000** per year*

Defer payments until after graduation.**
Fast preliminary approval, usually in minutes.

 **Apply Now!**
Go to gmacbankfunding.com

Apply online in as little as 15 minutes

- Loans up to \$40,000 per academic year*
- Good for tuition and other educational expenses: books, fees, a laptop, room and board, travel home, etc.
- Get a check in as few as 5 business days
- Start payments now or up to six months after graduation**
- Reduce your interest rate by as much as 0.50% with automatic payments***

All loans are subject to application and credit approval.

* Undergraduate and graduate borrowers may borrow annually up to the lesser of the cost of attendance or \$30,000 (\$40,000 for certain schools where it has been determined that the annual cost of attendance exceeds \$30,000). Borrowers in the Continuing Education loan program may borrow annually up to \$30,000.

** Undergraduate students may choose to defer repayment until six months after graduation or ceasing to be enrolled at least half time in school. Interest only and immediate repayment options also available.

*** A 0.25% interest rate reduction is available for borrowers who elect to have monthly principal and interest payments transferred electronically from a savings or checking account. The interest rate reduction will begin when automatic principal and interest payments start, and will remain in effect as long as automatic payments continue without interruption. This reduced interest rate will return to contract rate if automatic payments are cancelled, rejected or returned for any reason. Upon request, borrowers are also entitled to an additional 0.25% interest rate reduction if (1) the first 36 payments of principal and interest are paid on time, and (2) at any time prior to the 36th on time payment, the borrower who receives the monthly bill elects to have monthly principal and interest payments transferred electronically from a savings or checking account, and continues to make such automatic payments through the 36th payment. This reduced interest rate will not be returned to contract rate if, after receiving the benefit, the borrower discontinues automatic electronic payment. The lender and servicer reserve the right to modify or discontinue borrower benefit programs (other than the co-signer release benefit) at any time without notice.

GMAC Bank Member FDIC

Contents

20 Point Learning System	xi
Preface	xvii
Study Tips	xxiii

Chapter 1 ■ Operations with real numbers



1-1 Operations with fractions	2
1-2 Operations with decimals and percents	14
1-3 The set of real numbers and the real number line	25
1-4 Addition of real numbers	34
1-5 Subtraction of real numbers	41
1-6 Multiplication of real numbers	46
1-7 Division of real numbers	51
1-8 Properties of real numbers and order of operations	56
Chapter 1 lead-in problem	62
Chapter 1 summary	63
Chapter 1 error analysis	64
Chapter 1 critical thinking	64
Chapter 1 review	64


Chapter 2 ■ Solving equations and inequalities



2-1 Algebraic notation and terminology	67
2-2 Evaluating algebraic expressions	72
2-3 Algebraic addition and subtraction	79
2-4 The addition and subtraction property of equality	86
2-5 The multiplication and division property of equality	93
2-6 Solving linear equations	98
2-7 Solving literal equations and formulas	104
2-8 Word problems	107
2-9 Solving linear inequalities	113
Chapter 2 lead-in problem	122
Chapter 2 summary	122
Chapter 2 error analysis	123
Chapter 2 critical thinking	123
Chapter 2 review	124
Chapter 2 cumulative test	125

Chapter 3 ■ Polynomials and exponents

3-1	Exponents—I	127
3-2	Products of algebraic expressions	133
3-3	Exponents—II	139
3-4	Exponents—III	145
3-5	Scientific notation	148
	Chapter 3 lead-in problem	151
	Chapter 3 summary	152
	Chapter 3 error analysis	152
	Chapter 3 critical thinking	152
	Chapter 3 review	153
	Chapter 3 cumulative test	154

Chapter 4 ■ Factoring and solution of quadratic equations by factoring

4-1	Common factors	155
4-2	Factoring trinomials of the form $x^2 + bx + c$	162
4-3	Factoring trinomials of the form $ax^2 + bx + c$	166
4-4	Factoring the difference of two squares and perfect square trinomials	175
4-5	Other types of factoring	179
4-6	Factoring: A general strategy	184
4-7	Solving quadratic equations by factoring	186
4-8	Applications of the quadratic equation	193
	Chapter 4 lead-in problem	197
	Chapter 4 summary	198
	Chapter 4 error analysis	198
	Chapter 4 critical thinking	198
	Chapter 4 review	199
	Chapter 4 cumulative test	200

Chapter 5 ■ Rational Expressions, Ratio and Proportion

5-1	Rational numbers and rational expressions	202
5-2	Simplifying rational expressions	207
5-3	The quotient of two polynomials	212
5-4	Ratio and proportion	219
	Chapter 5 lead-in problem	228
	Chapter 5 summary	228
	Chapter 5 error analysis	228
	Chapter 5 critical thinking	229
	Chapter 5 review	229
	Chapter 5 cumulative test	230

Chapter 6 ■ Operations with Rational Expressions



6-1	Multiplication and division of rational expressions	232
6-2	Addition and subtraction of rational expressions	239
6-3	Addition and subtraction of rational expressions	245
6-4	Complex fractions	253
6-5	Rational equations	258
6-6	Rational expression applications	265
	Chapter 6 lead-in problem	270
	Chapter 6 summary	271
	Chapter 6 error analysis	271
	Chapter 6 critical thinking	272
	Chapter 6 review	272
	Chapter 6 cumulative test	274

Chapter 7 ■ Linear Equations in Two Variables




7-1	Ordered pairs and the rectangular coordinate system	276
7-2	Graphs of linear equations	289
7-3	The slope of a line	297
7-4	The equation of a line	305
7-5	Graphing linear inequalities in two variables	315
7-6	Functions defined by linear equations in two variables	323
	Chapter 7 lead-in problem	329
	Chapter 7 summary	329
	Chapter 7 error analysis	330
	Chapter 7 critical thinking	330
	Chapter 7 review	331
	Chapter 7 cumulative test	333

Chapter 8 ■ Systems of Linear Equations




8-1	Solutions of systems of linear equations by graphing	335
8-2	Solutions of systems of linear equations by elimination	340
8-3	Solutions of systems of linear equations by substitution	346
8-4	Applications of systems of linear equations	351
8-5	Solving systems of linear inequalities by graphing	360
	Chapter 8 lead-in problem	363
	Chapter 8 summary	364
	Chapter 8 error analysis	364
	Chapter 8 critical thinking	364
	Chapter 8 review	365
	Chapter 8 cumulative test	366

Chapter 9 ■ Roots and Radicals



9-1	Principal roots	367
9-2	Product property for radicals	373
9-3	Quotient property for radicals	377
9-4	Sums and differences of radicals	383
9-5	Further operations with radicals	386
9-6	Fractional exponents	391
9-7	Equations involving radicals	395
	Chapter 9 lead-in problem	400
	Chapter 9 summary	400
	Chapter 9 error analysis	401
	Chapter 9 critical thinking	401
	Chapter 9 review	401
	Chapter 9 cumulative test	402

Chapter 10 ■ Solutions of Quadratic Equations



10-1	Solutions of quadratic equations by extracting the roots	404
10-2	Solutions of quadratic equations by completing the square	409
10-3	Solutions of quadratic equations by the quadratic formula	416
10-4	Complex solutions to quadratic equations	424
10-5	The graphs of quadratic equations in two variables—quadratic functions	432
	Chapter 10 lead-in problem	444
	Chapter 10 summary	445
	Chapter 10 error analysis	445
	Chapter 10 critical thinking	446
	Chapter 10 review	446
	Final examination	447
	Appendix Answers and Solutions	449
	Index	505

Index

A

Abscissa of a point, 281
 Absolute value, 31
 Addition
 of decimals, 16
 of fractions, 7–9, 239
 of rational expressions, 239, 246
 Addition, identity element of, 35
 Addition, properties of
 associative, 38
 commutative, 35
 identity, 35
 rational expressions, 239, 246
 Addition and subtraction property of equality, 88, 340
 Addition and subtraction property of inequalities, 115
 Addition of
 algebraic expressions, 80–81
 like radicals, 384–85
 more than two real numbers, 42
 rational expressions, 239, 246
 two negative numbers, 36
 two numbers with different signs, 36–37
 two positive numbers, 34
 two real numbers, 38
 Additive inverse, 37–38
 Algebraic expression, 67
 Approximately equal to, 19, 29, 369
 Associative property of
 addition, 38
 multiplication, 49
 Axes, 280
 Axis of symmetry, 437

B

Base, 56, 127
 Binomial, 68
 Binomial, square of, 135, 410
 Boundary line, 316
 Braces, 25, 42
 Brackets, 42

C

Cantor, Georg, 25
 Coefficient, 68
 Common denominator, 7
 Commutative property of
 addition, 35
 multiplication, 46
 Completely factored form, 157

Completing the square, 411–13
 Complex fractions, 253
 primary denominator, 253
 primary numerator, 253
 secondary denominators of, 253
 simplifying, 253–55
 Complex numbers, 425
 addition of, 425
 multiplication of, 426
 rationalizing the denominator of, 427
 Components of an ordered pair, 277
 Compound inequality, 114
 Conjugate factors, 388
 Consistent system of equations, 337
 Constant, 67
 Coordinate, 29, 281
 Coordinates of a point, 281
 Counting numbers, 26

D

Decimal number, 14
 fraction, 14
 point, 14
 repeating, 19, 27
 terminating, 27
 Decreasing, 29
 Degree of a polynomial, 69, 186
 Denominator
 of a fraction, 2
 least common, 42
 of a rational expression, 203
 Denominator, rationalizing a, 427, 480, 482
 Dependent system, 337
 Descending powers, 69, 163
 Difference of two squares, 136
 Discriminant, 429
 Distance-rate-time, 266, 352
 Distributive property of
 multiplication over addition, 79
 Dividend, 51
 Division
 monomial by monomial, 212
 polynomial by monomial, 212
 polynomial by polynomial, 213–16
 property of rational expressions, 234
 of rational expressions, 235
 Division, definition of, 51
 Division by zero, 53
 Division involving zero, 53
 Division of
 decimals, 17
 fractions, 5, 235
 like bases, 140
 two or more real numbers, 52
 two real numbers, 52

Divisor, 51
 Domain
 of a rational expression, 204–5
 of a relation, 323

E

Element of a set, 25
 Elimination, method of solving systems of linear equations, 340–44
 Empty set, 260, 398
 Equality, properties of
 addition and subtraction, 88
 multiplication and division, 94
 squaring, 396
 symmetric, 89
 Equation, parts of an, 86
 Equation of a line, 305
 Equations
 conditional, 87
 diagram of, 86
 equivalent, 87
 first-degree, 87
 graph of, 282–83, 289–94
 identical, 87
 linear, 87, 276
 literal, 104
 quadratic, 186, 404–31
 radical, 395–96
 rational, 258
 system of, 335
 Equivalent rational expressions, 245–46
 Evaluating rational expressions, 203–4
 Evaluation, 72
 Expanded form, 56, 127
 Exponent, 56, 127
 Exponential form, 56, 127
 Exponents, properties and definitions of
 definition, 127
 fraction, 139
 fractional, 392
 group of factors to a power, 129
 negative exponents, 141
 power of a power, 130
 product, 128
 quotient, 140
 zero as an exponent, 142
 Extracting the roots, 405–6
 Extraneous solutions, 260, 396

F

Factor, 3, 46
 Factor, greatest common, 156
 Factored form, completely, 157

Factoring

- common factors, 156
- difference of two cubes, 179–80
- difference of two squares 175–76
- four-term polynomials, 159
- by inspection, 170–74
- perfect square trinomials, 177–78
- strategy, 184
- sum of two cubes, 181–82
- trinomials, 162–64, 166–74

FOIL, 135**Formulas, 74, 104****Four-term polynomials, 159****Fraction, 2**

- complex, 253
- improper, 2
- proper, 2

Fraction exponents, 392**Fraction to a power, 139****Function, 324**

- domain of a, 325
- range of a, 325

Fundamental principle of rational expressions, 207**G****Graph**

- of a linear equation, in two variables, 282–83, 289–94
- of a point, 29, 359
- a quadratic equation, 432–40
- of systems of equations, 336–37

Graphing

- a linear equation in two variables, 289–94
- linear inequalities in two variables, 315–19
- quadratic functions, 434–39
- systems of linear equations, 336–37
- systems of linear inequalities, 360–61

Greatest common factor, 155–56**Grouping symbols, 42, 82****Grouping symbols, removing, 82****Group of factors to a power, 129****H****Half-planes, 316****Horizontal line, 292–93**

- slope of, 300

I**Identical equation, 87****Identity, 87****Identity element**

- of addition, 35
- of multiplication, 47

Inconsistent system, 337**Increasing, 29****Independent system, 337****Indeterminate, 53****Index, 370****Inequalities**

- compound, 114

linear, 113

- strict, 30
- in two variables, 315–19
- weak, 30

Inequalities, properties of

- addition and subtraction, 115
- multiplication and division, 115

Inequality symbols, 30, 113**Integers, 26****Intercepts, x- and y-, 290****Inverse property**

- additive, 38
- multiplicative, 94

Irrational numbers, 27, 368**L****Least common denominator (LCD), 7, 242****Least common multiple (LCM), 242****Like bases, 128****Like radicals, 383****Like terms, 80****Linear equation, 87****Linear equations in two variables, 276**

- graphs of, 289–93
- systems of, 335

Linear inequalities in two variables, 315

- graphs of, 315–19

Linear inequality, 113**Lines**

- equation of, 305
- horizontal, 300
- parallel, 309
- perpendicular, 310
- point-slope form, 306
- slope-intercept form of, 307
- slope of, 297–302
- standard form of, 305
- vertical, 301

Literal equation, 104**M****Mathematical statement, 86****Member of an equation, 86****Member of a set, 25****Mixed number, 5****Monomial, 68****Multinomial, 69****Multiple, least common, 242****Multiplication**

- of decimals, 16
- of fractions, 4
- identity element of, 47
- of rational expressions, 233
- symbols for, 46

Multiplication, properties of

- associative, 49
- commutative, 46
- fractions, 4, 232
- identity, 47

- rational expressions, 233

Multiplication and division property for inequalities, 115**Multiplication and division property of equality, 94****Multiplication of**

- like bases, 128
- monomial and a multinomial, 133
- monomials, 130
- multinomials, 134
- n th roots, 374
- square roots, 373
- two negative numbers, 47
- two numbers with different signs, 47
- two or more real numbers, 48
- two positive numbers, 46
- two real numbers, 48

Multiplicative inverse, 94**N****Natural numbers, 26****Negative exponents, 141****Negative integers, 26****Negative of, 31** **n th root**

- addition of, 383–85
- definition of, 370
- of a fraction, 378, 380
- rationalizing, 379–81, 388

Number line, 29**Numerator, 2, 203****Numerical coefficient, 68****O****Open sentence, 86****Opposite of, 31****Ordered pairs, first component of, 277****Ordered pairs, second component of, 277****Ordered pairs of numbers, 277****Order of operations, 57****Order relationships, 30****Ordinate of a point, 281****Origin, 29, 280****of a rectangular coordinate plane, 280****P****Parabola, 434**

- axis of symmetry of a, 437
- graphing a, 434–39
- intercepts of a, 435
- vertex of a, 436–37

Parallel lines, 309**Parentheses, 42****Percent, 19–20****Percentage, 20–21****Perfect cube, 180****Perfect square, 175–76****Perfect square integer, 368****Perfect square trinomial, 135, 409****Perimeter, 10, 74****Perpendicular lines, 310****Pi (π), 28****Point**

- abscissa of a, 281
- graph of a, 281
- ordinate of a, 281

Point-slope form, 306**Polynomial, 68**

Polynomial, degree of, 69, 186
 Positive integers, 26
 Power of a power, 130
 Prime factor form, 3
 Prime number, 3
 Prime polynomial, 164
 Principal n th root, 370
 Principal square root, 368
 Product, 3, 46
 Product of rational expressions, 233
 Product property of exponents, 128
 Product property of radicals, 374
 Product property of square roots, 373
 Properties of real numbers, 56
 Proportion, 221
 property of, 221
 terms of, 221
 Pythagorean Theorem, 369

Q

Quadrants, 280
 Quadratic
 equations, 186, 404–31
 formula, 417–19
 standard form of, 187, 404
 Quotient, 5, 51
 of two polynomials, 212–16
 Quotient property of exponents, 140
 Quotient property of radicals, 378, 380

R

Radical equation, 395–96
 Radicals, properties of
 product, 373, 374
 quotient, 378, 380
 Radical symbol, 368
 Radicand, 368
 Range of a relation, 323
 Ratio, 219
 Rational equation, 258
 application, 265–67
 in more than one variable, 261
 Rational expression, 202
 Rational expressions
 applications of, 265–67
 completely reduced, 208–10
 definition of, 202
 denominator of a, 203
 difference of, 239, 246
 domain of, 204
 fundamental principle of, 207
 least common denominator of, 242
 numerator of, 203
 product of, 233
 quotient of, 235
 sum of, 239, 246
 Rationalizing a denominator, 379–81, 388, 427
 Rational number, 27
 Real number, 28
 Real number line, 29
 Reciprocal, 5, 94
 Rectangular coordinate plane, 280
 Reducing to lowest terms, 4, 208

Relation, 323
 domain of a, 323
 range of a, 323
 Remainder, 5
 Repeating decimal, 19
 Root, 86
 Root, extraneous, 260

S

Scientific notation, 148
 computation using, 149
 standard form of, 149
 Set, 25
 Set symbolism, 25
 Signed numbers, 34
 Similar terms, 80
 Simple interest, 112
 Simplifying rational expressions, 207
 Slope, 297–98
 definition of, 298–99
 of a horizontal line, 300
 of parallel lines, 309
 of perpendicular lines, 310
 undefined, 300–1
 of a vertical line, 301
 Slope-intercept form, 307
 Solution, 86, 276
 extraneous, 260
 Solution set, 86, 187
 of a linear equation in two variables, 277
 Solutions of quadratic equations
 by completing the square, 411–13
 complex solutions of, 427
 by extracting the roots, 405–6
 by factoring, 404–5
 by quadratic formula, 418–19
 Special products, 135–37
 Square of a binomial, 135
 Square root, 367
 Square root property, 405
 Standard form of a quadratic equation, 404
 Standard form of the equation of a line, 305
 Statement, mathematical, 86
 Straight line, 289
 Strict inequality, 30
 Subscripts, 75
 Subset, 26
 Substitution property, 72
 method of solving a system of linear equations, 346–48
 Subtraction, definition of, 41
 decimals, 16
 Subtraction, of rational expressions, 239
 Subtraction of
 fractions, 7–9, 239
 more than two real numbers, 42
 two real numbers, 41
 Symbols
 absolute value, 31
 braces, 25, 42
 brackets, 42
 is an element of, 26
 is approximately equal to, 19
 is a subset of, 26
 is greater than, 30
 is greater than or equal to, 30

is less than, 30
 is less than or equal to, 30
 multiplication dot, 46
 null set, 260
 parentheses, 42
 Symmetric property of equality, 89
 Symmetry, 31
 Systems of linear equations, 335
 applications of, 351–55
 consistent and independent, 337–38
 dependent, 337–38
 graphing of, 336–37
 inconsistent, 337–38
 linear, 335
 solution by elimination, 340–44
 solution by substitution, 346–48
 Systems of linear inequalities, 360–61

T

Term, 67
 Trinomial, 68
 Trinomial, perfect square, 135, 409

U

Undefined, 53
 Undefined slope, 301
 Undirected distance, 31

V

Variable, 29, 67
 Vertex of a parabola, 436–37
 Vertical line, 293
 slope of, 301

W

Weak inequality, 30
 Whole numbers, 26

X

x -axis, 280
 x -intercept, 290, 435

Y

y -axis, 280
 y -intercept, 290, 435

Z

Zero, division involving, 53
 Zero as an exponent, 142
 Zero factor property, 47
 Zero product property, 187

Campfire queen Cycling champion Sentimental geologist*

Learn more about
Marjon Walrod
and tell us more
about you. Visit
pwc.com/bringit.

Your life. You can
bring it with you.



*connectedthinking

PRICEWATERHOUSECOOPERS 

© 2006 PricewaterhouseCoopers LLP. All rights reserved. "PricewaterhouseCoopers" refers to PricewaterhouseCoopers LLP (a Delaware limited liability partnership) or, as the context requires, the PricewaterhouseCoopers global network or other member firms of the network, each of which is a separate and independent legal entity. *connectedthinking is a trademark of PricewaterhouseCoopers LLP (US). We are proud to be an Affirmative Action and Equal Opportunity Employer.